arXiv:hep-th/9407057v1 11 Jul 1994

Integrable highest weight modules over affine superalgebras and number theory

by Victor G. Kac[1*] and Minoru Wakimoto[2]

Department of Mathematics, M.I.T., Cambridge, MA 02139

Department of Mathematics, Mie University, Tsu 514, Japan

To Bertram Kostant from whom we learned so much.

§0. Introduction.

The problem of representing an integer as a sum of squares of integers has had a long history. One of the first after antiquity was A. Girard who in 1632 conjectured that an odd prime p can be represented as a sum of two squares iff $p \equiv 1 \bmod 4$, and P. Fermat in 1641 gave an "irrefutable proof" of this conjecture. The subsequent work on this problem culminated in papers by A.M. Legendre (1798) and C.F. Gauss (1801) who found explicit formulas for the number of representations of an integer as a sum of two squares. C.G. Bachet in 1621 conjectured that any positive integer can be represented as a sum of four squares of integers, and it took efforts of many mathematicians for about 150 years before J.-L. Lagrange gave a proof of this conjecture in 1770.

A radically new approach to the problem, based on the theory of elliptic functions, was found by C.G.J. Jacobi in his famous *Fundamenta nova* [J] published in 1829. In order to state his results, introduce the following power series in q:

$$\Box(q) = \sum_{j \in \mathbb{Z}} q^{j^2}.$$

Then, clearly, the coefficient of q^k in the power series expansion of $\Box(q)^d$ is the number of representations of k as a sum of d squares of integers (taking into account the order of summands). Jacobi discovered the following remarkable identities, which gave a solution to the problem for $d = 2$, 4, 6 and 8:

$$(0.1) \qquad \Box(q)^2 = 1 + 4 \sum_{j,k=1}^{\infty} (-1)^k q^{j(2k-1)},$$

---

[*] Supported in part by NSF grant DMS-9103792.

$$\square(q)^4 = 1 + 8 \sum_{\substack{j,k=1 \\ 4 \nmid k}}^{\infty} k q^{jk}, \tag{0.2}$$

$$\square(q)^6 = 1 - 4 \sum_{\substack{j,k=1 \\ j>k, j-k \,\text{odd}}}^{\infty} (-1)^{\frac{1}{2}(j-1)(k+1)} (j^2 - k^2) q^{\frac{1}{2}jk}, \tag{0.3}$$

$$\square(q)^8 = 1 + 16 \sum_{j,k=1}^{\infty} (-1)^{(j+1)k} k^3 q^{jk}. \tag{0.4}$$

Further development has been based, starting with the works of Jacobi and Legendre, on the theory of modular forms. Explicit formulas for odd $d \leq 7$ have been found by Legendre, Eisenstein, Smith, Hardy and Ramanujan (the case of odd d is much harder than that of even d). In the 1860s J. Liouville obtained a series of beautiful formulas for $d = 10$ and 12. In 1917 L.J. Mordell showed how in principle the problem can be solved for even d using the Fricke-Klein theory, but the difficulties increase rapidly with d. One can find a detailed (but quite disordered) account on sums of squares in Volume II of Dickson's "History of the Theory of Numbers" (Chapters VI–IX).

The related problem of representing a positive integer by sums of triangular numbers (*i.e.,* numbers of the form $\frac{1}{2}j(j+1)$, $j = 0, 1, \ldots$) has also been quite popular, and a number of deep results was obtained by Gauss, Legendre and others (see Chapter 1 of Vol. 2 of Dickson's "History"). Here the problem, clearly, is to find a power series expansion of powers of the series

$$\Delta(q) = \sum_{j=0}^{\infty} q^{\frac{1}{2}j(j+1)}.$$

The main objective of the present paper is to demonstrate that representation theory of affine superalgebras is a natural framework for the study of the problem of sums of squares and sums of triangular numbers. However, in order to state our main result we need only the following notions. Let V be a finite-dimensional vector space over $\mathbb{R}$ with a non-degenerate symmetric bilinear form $(.|.)$. Let Δ be a finite set of vectors in V not containing 0, called the set of *roots*, and let $\Delta = \Delta_0 \cup \Delta_1$ be represented as a disjoint union of two subsets, called the sets of *even* and *odd roots*. A choice of a linear function f on V which does not vanish on Δ defines a subset of *positive roots* $\Delta_+ = \{\alpha \in \Delta | f(\alpha) > 0\}$; we let $\Delta_{\epsilon +} = \Delta_\epsilon \cap \Delta_+$, $\epsilon = 0, 1$. Given $\alpha \in V$ we define the exponential function e^α

by $e^\alpha(v) = e^{(\alpha|v)}$. We call the *Weyl denominator* associated to the above *root data* the function

$$\mathrm{R} = \prod_{\alpha\in\Delta_{0+}} (1-e^{-\alpha}) \Big/ \prod_{\alpha\in\Delta_{1+}} (1+e^{-\alpha}).$$

Let q be an indeterminate; we call the *affine denominator* the infinite product

$$\hat{\mathrm{R}} = \mathrm{R} \prod_{n=1}^{\infty} \left((1-q^n)^\ell \prod_{\alpha\in\Delta_0} (1-q^n e^\alpha) \Big/ \prod_{\alpha\in\Delta_1} (1+q^n e^\alpha) \right),$$

where ℓ is the dimension of the $\mathbb{R}$-span of Δ. Our main objective is to find expansions for R and $\hat{\mathrm{R}}$. Of course, in the case when $\Delta = \Delta_0$ is the set of roots of a simple finite-dimensional Lie algebra, an expansion for R (resp. $\hat{\mathrm{R}}$) is given by the well-known Weyl denominator identity (resp. Macdonald identity [Mac]), see *e.g.,* [K9]. Our main Theorems 2.1 and 4.1 extend these results to the case when Δ is a set of roots of a finite- dimensional simple Lie superalgebra $\mathfrak{g}$ with a non-degenerate Killing form.

The connection to sums of squares is obvious. Indeed

$$\frac{\widehat{\mathrm{R}}}{\mathrm{R}}(0) = \prod_{n=1}^{\infty} (1-q^n)^{\ell+|\Delta_0|} \Big/ (1+q^n)^{|\Delta_1|},$$

and if it happens that

$$\ell + |\Delta_0| = |\Delta_1| =: d, \tag{0.5}$$

then we get an expansion of $\square(q)^d$, in view of Gauss' identity

$$\square(q) = \prod_{n=1}^{\infty} \frac{1-q^n}{1+q^n}.$$

Evaluating at points other than 0, one gets expansions for triangular numbers (and, more generally, m-gonal numbers).

To be more specific, let us give an example of a root system of type $A(m-1, n-1)$, where $m > n \geq 1$. Let V be the $(m+n)$-dimensional vector space over $\mathbb{R}$ with basis $\epsilon_1, \ldots, \epsilon_m, \delta_1, \ldots, \delta_n$ and the symmetric bilinear form given by $(\epsilon_i|\epsilon_i) = -(\delta_j|\delta_j) = 1$ for all i and j and zero for any other pair. Let $\Delta_0 = \{\epsilon_i - \epsilon_j | 1 \leq i, j \leq m, i \neq j\} \cup \{\delta_i - \delta_j | 1 \leq i, j \leq n, i \neq j\}$, $\Delta_1 = \{\epsilon_i - \delta_j, \delta_j - \epsilon_i | 1 \leq i \leq m, 1 \leq j \leq n\}$, $\Delta = \Delta_0 \cup \Delta_1$. Let

$\Delta_+ = \{\epsilon_i - \epsilon_j, \delta_i - \delta_j | i < j\} \cup \{\epsilon_i - \delta_j | i \leq j\} \cup \{\delta_i - \epsilon_j | i < j\}$. Condition (0.5) holds iff $m = n+1$. In this case, *i.e.,* Δ of type $A(n, n-1)$, the denominator identity is

$$\tag{0.6} \begin{aligned} \mathrm{R}_n := \prod_{i<j} \left(1 - \frac{x_i}{x_j}\right) \prod_{i<j} \left(1 - \frac{y_i}{y_j}\right) \prod_{i \leq j} \left(1 + \frac{x_i}{y_j}\right)^{-1} \prod_{i<j} \left(1 + \frac{y_i}{x_j}\right)^{-1} \\ = \sum_{w \in S_{n+1}} \epsilon(w) \prod_{i=1}^{n} \left(1 + \frac{y_i}{x_{w(i)}}\right)^{-1}. \end{aligned}$$

Here $x_i = e^{-\epsilon_i}$, $y_i = e^{-\delta_i}$, i and j take all possible values and $\epsilon(w)$ is the sign of the permutation $w \in S_{n+1}$. Furthermore, the affine denominator identity reads:

(0.7)

$$\begin{aligned} \mathrm{R}_n \prod_{k=1}^{\infty} \left((1-q^k)^{-1} \prod_{\substack{1 \leq i,j \leq n+1 \\ 1 \leq r,s \leq n}} \left(1 - q^k \frac{x_i}{x_j}\right) \left(1 - q^k \frac{y_r}{y_s}\right) \Big/ \left(1 + q^k \frac{x_i}{y_r}\right) \left(1 + q^k \frac{y_s}{x_j}\right) \right) \\ = \sum_{w \in S_{n+1}} \epsilon(w) \sum_{\vec{m} \in M_n} q^{\frac{1}{2} \sum_{i=1}^{n+1} m_i^2} \vec{x}^{\vec{m}} \prod_{i=1}^{n} \left(1 + \frac{y_i}{x_{\sigma(i)}} q^{-m_i}\right)^{-1}. \end{aligned}$$

Here M_n is the sublattice of the lattice $\sum_i \mathbb{Z}\epsilon_i$ of vectors with zero sum of coordinates (= the root lattice of type A_n) and $\vec{x}^{\vec{m}} := x_1^{m_1} \dots x_{n+1}^{m_{n+1}}$. Dividing both sides of (0.7) by R and taking limit as x_i, $y_i \to 1$, we obtain an expansion of $\square(q)^{2n(n+1)}$. One of the ways to actually compute this limit is explained in the proof of Theorem 4.2, but probably there is a different way giving a better formula. Of course, for $n = 1$ we thus recover Jacobi's formula (0.2). Another specialization of (0.7) gives the following expansion (see Example 4.3b):

$$\tag{0.8} \square(-q)^{2n} = 2^n \sum_{\vec{m} \in M_n} \sum_{s \in \mathbb{S}} (-1)^{|s|} q^{\frac{n+1}{2} \sum_{i=1}^{n+1} (m_i^2 + m_i - m_i \tilde{s}_i)} \prod_{j=1}^{n} \left(1 + q^{(n+1)(m_j - \tilde{s}_j)}\right)^{-1},$$

where $\mathbb{S}$ is the set of all subsets of the set $\{\epsilon_i - \epsilon_j | i < j\}$ (= set of positive roots of type A_n), and $\tilde{s}$ (resp. $|s|$) denote the sum (resp. number) elements from $s \in \mathbb{S}$. Formula (0.8) for $n = 1$ is an equivalent form of (0.1).

Even the simplest case of (0.6), when $n = 1$, is a truly remarkable identity. Letting $u = -x_1/y_1$, $v = -x_2/y_1$, it takes the following beautiful symmetric form:

(0.9)

$$\prod_{n=1}^{\infty} \frac{(1-q^n)^2 (1 - uvq^{n-1})(1 - u^{-1}v^{-1}q^n)}{(1 - uq^{n-1})(1 - u^{-1}q^n)(1 - vq^{n-1})(1 - v^{-1}q^n)} = \left(\sum_{m,n=0}^{\infty} - \sum_{m,n=-1}^{-\infty} \right) u^m v^n q^{mn}.$$

After establishing this identity, we realized that it has been rediscovered many times in the past. It is implicitly contained in [KL] as it follows immediately by comparing characters in the super boson-fermion correspondence (which gives a combinatorial proof of (0.9)), and in [M] where it is used to rewrite the characters of $N = 2$ superconformal algebras in a multiplicative form. As has been pointed out in [F] and [H], (0.9) is a special case of the Ramanujan summation formula for the bilateral basic hypergeometric function ${}_1\Psi_1$.

Condition (0.5) holds iff $\mathfrak{g}$ is of type $A(n, n-1)$, $B(n, n)$ or $D(n+1, n+1)$ ($n = 1, 2, \ldots$), which gives an expansion for $\Box(q)^N$ for $N = 2n(n+1)$, $2n(2n+1)$ and $4(n+1)^2$ respectively (given by Theorem 4.2; see also Examples 5.3 and 5.4). The first members of the first and the second series are Jacobi's formulas (0.2) and (0.3). The second member of the first series gives an expansion for $\Box(q)^{12}$, however we do not know how to derive from this expansion the famous result of J. Liouville (1860) that the number of representations of $2k$, where k is odd, as a sum of 12 squares equals $264 \sum_{j|k} j^5$. Finally note that Jacobi's formula (0.4) is obtained by evaluating at 0 the (conjectured) denominator identity for the case $\mathfrak{g} = gl(2, 2)$, which is a Lie superalgebra with a zero Killing form (see Conjecture 7.1). Another very interesting case which is not covered by Theorem 4.1 is $\mathfrak{g} = Q(m)$ (see Conjecture 7.2). A specialization of the corresponding affine denominators gives very beautiful power series expansions of $\Delta(q)^{4s^2}$ and $\Delta(q)^{4s(s+1)}$ (see Section 7). For $s = 1$ we thus recover Legendre's formulas for sums of 4 and 8 triangular numbers (whereas formulas for 2, 4 and 6 triangular numbers are obtained by certain specializations of the affine denominator identities for $A(1, 0)$, $A(1, 0)$ and $B(1, 1)$ respectively, see Section 5). For $s \geq 2$ we obtain formulas which are probably new. For example, we find that the number of representations of n as a sum of 16 triangular numbers equals

$$\frac{1}{3 \cdot 4^3} \sum_{\substack{a,b,r,s \in \mathbb{N}\text{odd} \\ a > b \\ ar+bs=2n+4}} ab(a^2 - b^2)^2.$$

In Section 1 we recall the necessary material on a simple finite-dimensional Lie superalgebra $\mathfrak{g}$ [K2]. The new material here concerns the dual Coxeter number $h^\vee$ (which is the eigenvalue of the Casimir operator associated to the suitably normalized Killing form). As in the Lie algebra case, $h^\vee$ plays an important role in the theory of affine superalgebras. In Section 2 we introduce the important notion of the defect of $\mathfrak{g}$ (which is 0 in the Lie algebra case) and of a maximal isotropic set of roots. This allows us to state the denominator identity (Theorem 2.1) for the root system of $\mathfrak{g}$. We give a proof which works in the case of def $\mathfrak{g} \leq 1$ (which takes care of all exceptional Lie superalgebras). A proof in the general case involves more calculations and will be published elsewhere. At any rate, the proof is based on the analysis of irreducible subquotients of the Verma module $M(0)$ over $\mathfrak{g}$ with highest weight 0. Since the denominator identity may be viewed as the character of the 1-dimensional $\mathfrak{g}$-module, it is only natural to go on and try to compute in Section 3 the character of an arbitrary finite-dimensional irreducible $\mathfrak{g}$-module V. There has been a number of papers concerned with this problem [K3, K4, BL, TM1, TM2, VJ1, VJ2, JHKT1, JHKT2, P, PS1, PS2, S, Se, ...]. Our basic new ingredient is the notion of a λ-maximal isotropic subset, which allows us to define atypicality of V which is independent of the choice of the set of positive roots, and of a tame $\mathfrak{g}$-module V. Our first basic result is Theorem 3.1 which states that V is tame (and hence gives a formula for $\operatorname{ch} V$) in the case when for some choice of the set of positive roots there exists a $\Lambda+\rho$-maximal isotropic subset consisting of simple roots. (Note that Theorem 2.1 is a special case of Theorem 3.1 since def $\mathfrak{g} = \operatorname{atp} 0$.) This theorem covers most of the known formulas for $\operatorname{ch} V$ (proved in [K3, K4, BL, VJ1, VJ2, JHKT2, ...]), but unfortunately there exist V that are not tame. We describe all cases of non-tame V of atypicality ≤ 1; this gives, in particular, character formulas for all V over exceptional Lie superalgebras (Example 3.3). We also give a proof of a dimension formula (Theorem 3.2), which is based on a nice regularization procedure, and of a superdimension formula (Theorem 3.3), which is based on an interesting property of the root system of $\mathfrak{g}$ (Lemma 2.2d) and on Theorem 2.1.

In the second part of the paper (Sections 4–8) we turn to the associated to $\mathfrak{g}$ affine superalgebra $\hat{\mathfrak{g}}$. The main result is Theorem 4.1 which gives the affine denominator identity in the case when the Killing form is nondegenerate (equivalently, $h^\vee \neq 0$). The proof is again based on the analysis of irreducible subquotients of the Verma module over $\hat{\mathfrak{g}}$ with zero highest weight. (Of course, this is the basic idea of the proof of Macdonald identities given in [K1].) Examples 4.4 and 4.5 give a more convenient form for specializations of the affine denominator identity. Theorems 4.2 and 4.3 and Example 4.3b give explicit formulas for two specializations of this identity. These formulas are applied in Section 5 to derive expressions for the number of representations of an integer as sums of squares and sums of triangular numbers. In Section 6 we use the simplest affine denominator identity to study transformation properties of certain "degenerate" theta functions and modular forms and apply this to prove transformation properties of the $N=2$ superconformal algebras, *cf.*

[RY]. (These results may be extended to the general case by the same method.) In Section 7 we discuss the affine denominator identity for $\mathfrak{g} = gl(2,2)$, the simplest case when $h^\vee = 0$. Another extremely interesting case is $\mathfrak{g} = Q(m)$. We conjecture the corresponding affine denominator identity and derive formulas for the number of representations as sums of triangular numbers in terms of dimensions of irreducible representations of sl_{m+1}. Finally, in Section 8 we introduce the notion of an integrable $\hat{\mathfrak{g}}$-module, based on our experience with the 1-dimensional module. Theorem 8.1 gives a classification of irreducible integrable highest weight $\hat{\mathfrak{g}}$-modules $\hat{L}(\Lambda)$.

(As in the Lie algebra case, the level of such a module is a non-zero integer, but there may be infinitely many such modules of positive level.) It is an extremely interesting problem to calculate the characters of integrable $\hat{L}(\Lambda)$. A partial answer is given for $A(1,0)^\wedge$ (Example 8.1). Of course, in the case when $\operatorname{def}\mathfrak{g} = 0$ the answer is well known for both $\mathfrak{g}$ and $\hat{\mathfrak{g}}$ [K5].

Acknowledgment

We learned from V. Serganova the very useful method of odd reflections. Discussions with her and I. Penkov were very illuminating. We relied extensively on the (unpublished) tables of J. Thierry-Mieg [TM1] and computer calculations of J. Van der Jeugt of dimensions and superdimensions of irreducible $\mathfrak{g}$-modules. Discussions with them and R.C. King were very useful. We are very grateful to all these people.

The results of this work were reported at the conference on universal enveloping algebras in Paris in June 1992 and at the Kostant conference in May 1993.

§1. Dual Coxeter number.

Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a finite-dimensional simple Lie superalgebra over $\mathbb{C}$ with a non-degenerate even invariant bilinear form B. Recall that "even" (resp. "invariant") means that $B(\mathfrak{g}_0, \mathfrak{g}_1) = 0$ (resp. $B\big([a,b],c\big) = B\big(a,[b,c]\big), a,b,c \in \mathfrak{g}$) and that all such forms are proportional. Denote by $h^\vee_B$ the eigenvalue of the Casimir operator Ω_B associated to B [K2, p. 85] in the adjoint representation.

Proposition 1.1. *$h^\vee_B \neq 0$ if and only if the Killing form is non-degenerate.*

Choose a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}_0$, let $\Delta \in \mathfrak{h}^*$ be the set of roots, Δ_0 and Δ_1 the subsets of even and odd roots and let $W \subset GL(\mathfrak{h}^*)$ be the Weyl group (= the group generated by reflections r_α with respect to even roots α). Let $\mathfrak{g} = \mathfrak{h} \oplus \Big(\underset{\alpha\in\Delta}{\oplus} \mathfrak{g}_\alpha\Big)$ be the root space decomposition.

Recall that an odd root α is isotropic $\big($i.e. $B(\alpha,\alpha) = 0\big)$ iff 2α is not a root. We let

$$\bar{\Delta}_0 = \big\{\alpha \in \Delta_0 \bigm| \tfrac{1}{2}\alpha \notin \Delta\big\}, \quad \bar{\Delta}_1 = \big\{\alpha \in \Delta_1 \bigm| (\alpha|\alpha) = 0\big\}.$$

Then $\Delta_1 \setminus \bar{\Delta}_1 = \frac{1}{2}(\Delta_0 \backslash \bar{\Delta}_0)$ and $\bar{\Delta}_0$ is a root system.

Choose a set of positive roots Δ_+ in Δ, let $\Delta_{\epsilon+} = \Delta_\epsilon \cap \Delta_+$ ($\epsilon = 0$ or 1) and let $\mathfrak{n}_+ = \underset{\alpha\in\Delta_+}{\oplus} \mathfrak{g}_\alpha$. Let $\Pi = \{\alpha_1, \dots, \alpha_l\} \subset \Delta_+$ be the set of simple roots and let $\theta \in \Delta_+$ be the highest root. Let ρ_ϵ be the half of the sum of the roots from $\Delta_{\epsilon+}$ and and let $\rho = \rho_0 - \rho_1$. Then, $B(\rho, \alpha_i) = \frac{1}{2}B(\alpha_i, \alpha_i)$, $i = 1, \dots, l$. Furthermore one has

$$h^\vee_B = B(\rho,\rho) + \tfrac{1}{2}B(\theta,\theta). \tag{1.1}$$

The following generalization of Freudenthal-de Vries "strange formula" holds:

$$B(\rho,\rho) = \frac{h^\vee_B}{12}(\dim \mathfrak{g}_0 - \dim \mathfrak{g}_1). \tag{1.2}$$

Assuming that $h^\vee_B \neq 0$, let

$$\Delta^\sharp_0 = \big\{\alpha \in \Delta_0 \bigm| h^\vee_B B(\alpha,\alpha) > 0\big\}$$

(this is independent of the choice of B), and let $W^{\sharp}$ denote the subgroup of W generated by reflection r_{α} with respect to all $\alpha \in \Delta_0^{\sharp}$. Denote by $(.|.)$ the even invariant bilinear form on $\mathfrak{g}$ normalized by the condition

$$(\alpha|\alpha) = 2 \text{ for the longest root } \alpha \in \Delta_0^{\sharp}. \tag{1.3}$$

The corresponding to this form number $h^{\vee} = h^{\vee}_{(.|.)}$ is called the *dual Coxeter number* of the Lie superalgebra $\mathfrak{g}$. We let $h^{\vee} = 0$ if $h^{\vee}_B = 0$. We give below the number $h^{\vee}$ in all cases (excluding Lie algebra cases):

$$\begin{aligned}
A(m,n) &: \ h^{\vee} = |m-n|; \quad C(n) : h^{\vee} = n-1; \\
B(m,n) &: \ h^{\vee} = 2(m-n)-1 \text{ if } m > n, \quad = n-m+\tfrac{1}{2} \text{ if } m \leq n; \\
D(m,n) &: \ h^{\vee} = 2(m-n-1) \text{ if } m \geq n+1, \quad = n-m+1 \text{ if } m < n+1; \\
D(2,1;\alpha) &: \ h^{\vee} = 0; \quad F(4) : h^{\vee} = 3; \quad G(3) : h^{\vee} = 2.
\end{aligned}$$

Remarks 1.1. (a) Let $\kappa(\ .\ ,\ .\ )$ denote the Killing form, which we assume to be non-degenerate. Then $\Delta_0^{\sharp} = \left\{\alpha \in \Delta_0 \;\middle|\; \kappa(\alpha,\alpha) > 0\right\}$ is the root system of one of the simple components of $\mathfrak{g}_0$ and $h^{\vee} = \kappa(\alpha,\alpha)^{-1}$ for the longest $\alpha \in \Delta_0^{\sharp}$.

(b) $h^{\vee} = 0$ if and only if $\mathfrak{g}$ is of type $A(n,n)$, $D(n+1,n)$ or $D(2,1;\alpha)$ [ ]. In these cases we define $\Delta_0^{\sharp}$ as follows. The root system Δ_0 is a union of two orthogonal to each other root subsystem; namely one has respectively:

$$\Delta_0 = A_n \cup A_n, \quad D_{n+1} \cup C_n, \quad D_2 \cup C_1.$$

Then we let $\Delta_0^{\sharp}$ to be the first subset and $W^{\sharp}$ to be the subgroup of W generated by the r_{α} with $\alpha \in \Delta_0^{\sharp}$. We normalize $(.|.)$ by (1.3) as before.

(c) There are many W-inequivalent choices of Δ_+. However, there is a unique (up to an automorphism of the root lattice) choice of Δ_+ such that Π contains a unique odd root; we call this Δ_+ *standard*.

In order to prove the above statements, recall that [K2, p. 85] the eigenvalue of the Casimir operator Ω_B in a representation with highest weight Λ is equal to $B(\Lambda, \Lambda + 2\rho)$ and that when the Killing form κ is non-degenerate one has:

$$\kappa(\theta, \theta + 2\rho) = 1 \tag{1.4}$$

provided that $\dim \mathfrak{g}_0 \neq \dim \mathfrak{g}_1$. Since

$$\dim \mathfrak{g}_0 = \dim \mathfrak{g}_1 \quad \text{iff} \quad \mathfrak{g} \simeq A(n+1, n),\ B(n,n) \text{ or } D(n,n) \tag{1.5}$$

and since (1.4) is a polynomial identity which holds for all other values of (m, n) (except when $\kappa \equiv 0$) we obtain that (1.4) holds in all cases when κ is non-degenerate. This proves Proposition 1.1 and formula (1.1).

From formula (2.1) in §2, it follows that

$$B(\rho, \rho) \text{ is independent of the choice of } \Delta_+. \tag{1.6}$$

Hence it suffices to check (1.2) for one choice of Δ_+. A more conceptual proof of (1.2) is based on the theory of "degenerate" theta functions (*cf.* Remark 6.1 in Section 6).

Example 1.1. $\mathfrak{g} = A(m,n)$ with $m > n$. Choose Δ_+ such that the number of odd simple roots is maximal. Then $\Pi = \Pi' \cup \Pi''$, where $(\alpha|\alpha) = 0$ if $\alpha \in \Pi'$ and $(\alpha|\alpha) = 2$ if $\alpha \in \Pi''$, and one can order the sets Π' and Π'':

$$\Pi' = \left\{\alpha_1, \alpha_2, \dots, \alpha_{2(n+1)}\right\}, \quad \Pi'' = \{\alpha_{2n+3}, \dots, \alpha_{m+n+1}\}$$

in such a way that all non-zero scalar products of distinct roots are:

$$(\alpha_{2j-1}|\alpha_{2j}) = 1 \text{ if } 1 \leq j \leq n+1, \quad (\alpha_{2j}|\alpha_{2j+1}) = -1 \text{ if } 1 \leq j \leq n,$$
$$(\alpha_j|\alpha_{j+1}) = -1 \text{ if } 2n+2 \leq j \leq m+n.$$

In particular, Π' (resp. Π'') is a set of simple roots of $A(n+1,n)$ (resp. A_{m-n-1}). The root system $\Delta_0^\sharp$ is of type A_m and has the following set of simple roots: $\{\alpha_1 + \alpha_2, \dots, \alpha_{2n+1} + \alpha_{2n+2}, \alpha_{2n+3}, \dots, \alpha_{m+n+1}\}$. In the (most important for us) case when $m = n+1$, we have $\Pi = \Pi'$ and $\Delta_0^\sharp$ has simple roots $\{\alpha_1 + \alpha_2, \dots, \alpha_{2n+1} + \alpha_{2n+2}\}$. Note that (1.2) for $A(m,n)$ follows from the "strange formula" for A_{m-n-1}.

§2. The Weyl denominator of $\mathfrak{g}$.

Given a subset of positive roots Δ_+ of Δ and a simple root α, we may construct a new subset of positive roots (*cf.* [PS2]):

$$\Delta'_+ = r_\alpha(\Delta_+) \ \text{ if } \ (\alpha|\alpha) \neq 0,$$
$$\Delta'_+ = \big(\Delta_+ \cup \{-\alpha\}\big) \setminus \{\alpha\} \ \text{ if } \ (\alpha|\alpha) = 0.$$

The set Δ'_+ is called a simple reflection of Δ_+. Then one has:

$$\rho' = \begin{cases} \rho - \alpha & \text{if} \quad (\alpha|\alpha) \neq 0 \\ \rho + \alpha & \text{if} \quad (\alpha|\alpha) = 0. \end{cases} \tag{2.1}$$

Note that Δ'_+ and Δ_+ are W-inequivalent if $(\alpha|\alpha) = 0$. By a usual argument, any two subsets of positive roots of Δ may be obtained from each other by a sequence of simple reflections.

By a *regular exponential function* on $\mathfrak{h}$ we mean a finite linear combination of exponentials e^λ, $\lambda \in \mathfrak{h}^*$. *A rational exponential* function is a ratio P/Q where P and Q are regular exponential functions and $Q \neq 0$. The Weyl group W acts on the field of rational exponential functions via $e^\lambda \mapsto e^{w(\lambda)}$, $w \in W$.

The rational exponential function

$$R = \prod_{\alpha \in \Delta_{0+}} (1 - e^{-\alpha}) \Big/ \prod_{\alpha \in \Delta_{1+}} (1 + e^{-\alpha}) \tag{2.2}$$

is called the *Weyl denominator* of $\mathfrak{g}$. Introduce also the *Weyl superdenominator*:

$$\begin{aligned} \check{R} &= \prod_{\alpha \in \Delta_{0+}} (1 - e^{-\alpha}) \Big/ \prod_{\alpha \in \Delta_{1+}} (1 - e^{-\alpha}) \\ &= \prod_{\alpha \in \bar{\Delta}_{0+}} (1 - e^{-\alpha}) \prod_{\alpha \in \Delta_{1+} \setminus \bar{\Delta}_{1+}} (1 + e^{-\alpha}) \Big/ \prod_{\alpha \in \bar{\Delta}_{1+}} (1 - e^{-\alpha}). \end{aligned} \tag{2.3}$$

The proof of the following lemma is straightforward.

Lemma 2.1. *Let Δ_+ and Δ'_+ be two subsets of positive roots in Δ, and let R and R' (resp. $\check{R}$ and $\check{R}'$) be the corresponding Weyl (resp. super) denominators. Then*

$$e^\rho R = (-1)^{|\Delta_{0+} \cap \Delta'_{0+}|} e^{\rho'} R', \quad e^\rho \check{R} = (-1)^{|\bar{\Delta}_{0+} \cap \bar{\Delta}'_{0+}|} e^{\rho'} \check{R}'.$$

The basic difference between the Lie algebra and superalgebra cases is that the restriction of the bilinear form $(.|.)$ to the subspace $V := \sum\limits_{\alpha\in\Delta} \mathbb{R}\alpha$ may be indefinite; the dimension of a maximal isotropic subspace of V is called the *defect* of $\mathfrak{g}$ and is denoted by $\operatorname{def}\mathfrak{g}$. Here is a list of defects

$$A(m-1,n-1),\ \ B(m,n),\ \ D(m,n)\ :\ \operatorname{def}\mathfrak{g} = \min(m,n)$$

$$C(n), D(2,1;\alpha), F(4), G(3)\ :\ \operatorname{def}\mathfrak{g} = 1.$$

(Note that $\operatorname{def}\mathfrak{g} = 0$ iff $\bar{\mathfrak{g}}$ is a Lie algebra or a Lie superalgebra $B(0,n)$.)

A subset S of Δ is called *maximal isotropic* if it consists of $\operatorname{def}\mathfrak{g}$ roots that span a maximal isotropic subspace of V. The following can be checked by a straightforward calculation.

Lemma 2.2. *(a) A maximal isotropic set of roots always exists.*

(b) Given a maximal isotropic subset of roots S there exists Δ_+ such that $S \subset \Pi$.

(c) Given two maximal isotropic subsets of roots $S = \{\beta_1, \dots, \beta_s\}$ and $S' = \{\beta'_1, \dots, \beta'_s\}$, there exists $w \in W^\sharp$ such that $w(\beta'_i) = \pm\beta_{\sigma(i)}$ for a permutation σ.

(d) Let $S \subset \bar{\Delta}_{1+}$ be a maximal isotropic subset and let $\bar{\Delta}^S_{0+} = \{\alpha \in \bar{\Delta}_{0+} | \alpha \perp S\}$. Then

$$|\bar{\Delta}_{1+} \setminus S| = |\bar{\Delta}_{0+} \setminus \bar{\Delta}^S_{0+}|.$$

Theorem 2.1. *Let S be a maximal isotropic subset of roots and choose Δ_+ such that $S \subset \Pi$. Then*

$$e^\rho R = \sum_{w \in W^\sharp} \epsilon(w) w \frac{e^\rho}{\prod_{\beta\in S}(1+e^{-\beta})} \tag{2.4}$$

This theorem is a special case of Theorem 3.1 in Section 3.

Here we shall give a proof in the case $\operatorname{def}\mathfrak{g} = 1$. The proof is based on the following

Lemma 2.3. *Suppose that an irreducible* $\mathfrak{g}$*-module with highest weight* $\lambda \in \mathfrak{h}^*$ *is a subquotient of a Verma* $\mathfrak{g}$*-module* $M(\Lambda)$*. Then there exists a sequence of weights* $\lambda_0 = \Lambda$, $\lambda_1, \dots, \lambda_k = \lambda$ *of* $M(\Lambda)$ *such that* λ_i $(0 < i \leq k)$ *is obtained from* λ_{i-1} *in one of the following two ways:*

(i) $\lambda_i = \lambda_{i-1} - n\gamma_i$*, where* $\gamma_i \in \Delta_{0+}$ *and* $(\lambda_{i-1} + \rho | \gamma_i^\vee) = n \in \mathbb{N}$*,*

(ii) $\lambda_i = \lambda_{i-1} - \gamma_i$*, where* $\gamma_i \in \bar{\Delta}_{1+}$ *and* $(\lambda_{i-1} + \rho | \gamma_i) = 0$*.*

Proof. See [K4, Theorem 3b]. (Note that the exponents in the determinant formula given by [K4, Theorem 3a] should be corrected as in [K8].) ■

Proof of Theorem 2.1 (case def $\mathfrak{g} = 1$.) As usual, we have:

$$e^\rho \mathrm{R} = \sum_{\lambda \in F} c_\lambda e^{\lambda + \rho}, \quad c_\lambda \in \mathbb{Z}, \quad c_\lambda \neq 0, \tag{2.5}$$

where F is the set of weights described by (i) and (ii) in Lemma 2.3 for $\Lambda = 0$. Choose Δ_+ such that Π contains a unique odd (isotropic) root, say, β. We clearly have:

$$\sum_{j \geq 0} c_{-j\beta} e^{\rho - j\beta} = \frac{e^\rho}{1 + e^{-\beta}}. \tag{2.6}$$

Consider the minimal i for which the possibility (ii) of Lemma 2.3 might occur. Then $\lambda_{i-1} = w(\rho) - \rho$, where $w = r_{\gamma_{i-1}} \dots r_{\gamma_1} \in W$ and $\lambda_i = \lambda_{i-1} - \gamma_i$, $\big(\rho | w^{-1}(\gamma_i)\big) = 0$, $\gamma_i \in \bar{\Delta}_1$. But by a case-wise inspection, $(\rho|\gamma) = 0$, $\gamma \in \bar{\Delta}_1 \Rightarrow \gamma = \pm\beta$. Hence $\gamma_i = w(\beta)$. Consider now the minimal $j \leq i-1$ for which $\gamma_j \notin \Delta_+^\sharp$, *i.e.*, $\gamma_j = \theta$. But $(\rho|\theta^\vee) < 0$, hence we conclude that $w \in W^\sharp$. Thus, $\lambda_i = w(\rho - \beta) - \rho$, $w \in W^\sharp$.

Furthermore, $(\beta|\theta^\vee) \geq 0$, hence $(\rho - \beta|\theta^\vee) < 0$, hence if s is minimal $> i$ for which possibility (ii) of Lemma 2.3 might occur, we conclude that $\lambda_{s-1} = w_1(\rho - \beta) - \rho$, $w_1 \in W^\sharp$.

By a case-wise inspection, we see that $(\rho - \beta|\beta_1) = 0$, $\beta_1 \in \bar{\Delta}_{1+} \Rightarrow \beta_1 = \beta$. Hence $\lambda_s = w_1(\rho - 2\beta)$, etc. Thus, the set $A := \{\lambda + \rho | \lambda \in F\}$ coincides with the set $\{w(\rho - j\beta) | w \in W^\sharp, j \in \mathbb{Z}_+\}$. But by Lemma 2.1, $w(e^\rho \mathrm{R}) = \epsilon(w) e^\rho \mathrm{R}$. This together with (2.6) completes the proof. ■

Remark 2.1. Recall that $\epsilon(w) = (-1)^m$, where m is the number of reflections with respect to roots from Δ_{0+} entering in a decomposition of w as a product of reflections. We let $\check{\varepsilon}(w) = (-1)^{\check{m}}$, where $\check{m}$ denote the number of reflections occurring in this decomposition with respect to roots from $\bar{\Delta}_{0+}$. Then one easily deduces from (2.4) the formula

$$e^\rho \check{R} = \sum_{w \in W^\sharp} \check{\epsilon}(w) w \frac{e^\rho}{\prod_{\beta \in S}(1 - e^{-\beta})}. \tag{2.7}$$

§3. Finite-dimensional irreducible $\mathfrak{g}$-modules.

Let $\lambda \in \mathfrak{h}^*$. The *atypicality* of λ, denoted by $\operatorname{atp}\lambda$, is the maximal number of linearly independent roots β_i such that $(\beta_i|\beta_j) = 0$ and $(\lambda|\beta_i) = 0$ for all i and j. Such a set $\{\beta_i\}$ is called a λ-*maximal isotropic* subset of Δ. (For example, a maximal isotropic subset $S \subset \Pi$ is a ρ-maximal isotropic subset.)

Proposition 3.1. *Given two λ-maximal isotropic subsets $\{\beta_i\}$ and $\{\beta'_i\}$ of Δ, there exists $w \in W$ such that $w(\lambda) = \lambda$ and $w(\beta_i) = \pm\beta'_{\sigma(i)}$ for some permutation σ.*

Let V be a finite-dimensional irreducible $\mathfrak{g}$ module. If we fix a set of positive roots Δ_+, we may talk about the highest weight Λ of V and about the ρ. If Δ'_+ is obtained from Δ_+ by a simple α- reflection, where $(\alpha|\alpha) = 0$, and if Λ' is the highest weight for Δ'_+, then we clearly have

$$\Lambda' = \Lambda - \alpha \quad \text{if} \quad (\Lambda|\alpha) \neq 0; \qquad \Lambda' = \Lambda \quad \text{if} \quad (\Lambda|\alpha) = 0. \tag{3.1}$$

Hence by (2.1) we obtain:

$$\begin{aligned} \Lambda' + \rho' &= \Lambda + \rho \quad \text{if} \quad (\Lambda + \rho|\alpha) \neq 0, \\ \Lambda' + \rho' &= \Lambda + \rho + \alpha \quad \text{if} \quad (\Lambda + \rho|\alpha) = 0. \end{aligned} \tag{3.2}$$

Corollary 3.1. *For the $\mathfrak{g}$-module V,* $\operatorname{atp}(\Lambda + \rho)$ *is independent of the choice of Δ_+.*

Thus, we may talk about the atypicality of the $\mathfrak{g}$-module V: $\operatorname{atp} V := \operatorname{atp}(\Lambda+\rho)$, which is independent of the choice of Δ_+. Note that $\operatorname{atp} V \leq \operatorname{def}\mathfrak{g}$.

Choose a $\Lambda+\rho$-maximal isotropic subset S_Λ in Δ_+ and consider the following rational exponential function:

$$\begin{aligned}\Sigma_\Lambda &:= \sum_{w\in W}\epsilon(w)w\Big(e^{\Lambda+\rho}\prod_{\beta\in S_\Lambda}\left(1+e^{-\beta}\right)^{-1}\Big),\\ \check{\Sigma}_\Lambda &:= \sum_{w\in W}\check{\epsilon}(w)w\Big(e^{\Lambda+\rho}\prod_{\beta\in S_\Lambda}\left(1-e^{-\beta}\right)^{-1}\Big).\end{aligned}$$

Denote by j_Λ the coefficient of $e^{\Lambda+\rho}$ in the regular exponential function $\prod_{\alpha\in\bar{\Delta}_{1+}}(1+e^{-\alpha})\sum_\Lambda$ and, provided that $j_\Lambda\neq 0$, let

$$\begin{aligned}\operatorname{ch}_\Lambda = j_\Lambda^{-1}e^{-\rho}R^{-1}\Sigma_\Lambda,\ \operatorname{sch}_\Lambda = j_\Lambda^{-1}e^{-\rho}\check{R}^{-1}\check{\Sigma}_\Lambda\\ \dim_\Lambda = \operatorname{ch}_\Lambda(0),\ \operatorname{sdim}_\Lambda = \operatorname{sch}_\Lambda(0).\end{aligned}$$

We shall write sometimes $\operatorname{ch}_{\Lambda,\Delta_+,S_\Lambda},\dots$ in place of $\operatorname{ch}_\Lambda,\dots$ to emphasize the dependence of Δ_+ and S_Λ.

Definition 3.1. *The $\mathfrak{g}$-module V is called tame with respect to a subset of positive roots Δ_+ and a subset $S_\Lambda\subset\Delta_+$, where Λ is the highest weight of V with respect to Δ_+ and S_Λ is a $\Lambda+\rho$-maximal isotropic subset of Δ_+, if $j_\Lambda\neq 0$ and the character of V is given by*

$$\operatorname{ch} V = \operatorname{ch}_{\Lambda,\Delta_+,S_\Lambda}\,. \tag{3.3}$$

Remarks 3.1. (a) Formula (3.3) implies

$$\operatorname{sch} V = \pm\operatorname{sch}_{\Lambda,\Delta_+,S_\Lambda}\,.$$

Here + or − occurs according as the highest weight vector is even or odd.

(b) If V is tame, then $\dim V = \dim_\Lambda, \operatorname{sdim} V = \pm\operatorname{sdim}_\Lambda$.

Remarks 3.2. (a) Suppose that Δ'_+ is obtained from Δ_+ by a simple α-reflection, where $(\alpha|\alpha)=0$. If $(\Lambda+\rho|\alpha)\neq 0$, then $\mathrm{ch}_{\Lambda,\Delta_+,S_\Lambda}=\mathrm{ch}_{\Lambda',\Delta'_+,S_\Lambda}$. If $(\Lambda+\rho|\alpha)=0$ and $\alpha\in\Pi\bigcap S_\Lambda$, we let $S'_\Lambda=(S_\Lambda\setminus\{\alpha\})\bigcup\{-\alpha\}$; then $\mathrm{ch}_{\Lambda,\Delta_+,S_\Lambda}=\mathrm{ch}_{\Lambda,\Delta'_+,S'_\Lambda}$. This follows from (3.1) and (3.2). However, if $(\Lambda+\rho|\alpha)=0$ and $\alpha\in\Pi\setminus S_\Lambda$, $\mathrm{ch}_{\Lambda',\Delta'_+,S_{\Lambda'}}$ may be different from $\mathrm{ch}_{\Lambda,\Delta_+,S_\Lambda}$.

(b) Let $S_\Lambda=\{\beta_1,\ldots,\beta_s\}$ and suppose $y\in W$ is such that $y(\Lambda+\rho)=\Lambda+\rho$, $y(\beta_i)=\beta_i$ for $i=2,\ldots,s$, and $y(\beta_1)=\epsilon(y)\beta'_1$, where $\beta'_1\in\Delta_+$. Let $S'_\Lambda=\{\beta'_1,\beta_2,\ldots,\beta_s\}$. Then $\mathrm{ch}_{\Lambda,\Delta_+,S_\Lambda}=\mathrm{ch}_{\Lambda,\Delta_+,S'_\Lambda}$.

(c) Theorem 2.1 states that the 1-dimensional $\mathfrak{g}$-module is tame with respect to (Δ_+,S_0) if $\Pi\supset S_0$. Note that then

$$j_0=\left|W/W^\sharp\right|. \tag{3.4}$$

The number j_0 is listed below:

$A(m,n)(m\geq n):(n+1)!$; $B(m,n):n!2^n$ if $m>n$, $2^m m!$ if $m\leq n$; $C(n):1$;
$D(m,n):n!2^n$ if $m\geq n+1$; $2^{m-1}m!$ if $m<n+1$; $D(2,1;\alpha)$, $F(4)$, $G(3):2$.

Proposition 3.2. *Let V be a finite-dimensional irreducible $\mathfrak{g}$-module. Choose a set of positive roots Δ_+ and let Λ denote the highest weight of V.*

(a) (cf. *[K4]) Let V be a typical $\mathfrak{g}$-module,* i.e., $(\Lambda+\rho|\alpha)\neq 0$ *for $\alpha\in\bar\Delta_1$ (equivalently:* $\mathrm{atp}\,V=0$*). Then V is tame with respect to $(\Delta_+,\emptyset)$.*

(b) (cf. *[BL], [JHKT2, VJ2]) If $\mathfrak{g}$ is of type $A(m,n)$ or $C(n)$ and* $\mathrm{atp}\,V=1$, *then V is tame with respect to (Δ_+,S_Λ) for any choice of S_Λ and $j_\Lambda=1$. (Note that* $\mathrm{atp}\,V=0$ *or* 1 *if $\mathfrak{g}$ is of type $A(m,0)$ or $C(n)$.)*

(c) If $(\Lambda+\rho|\alpha)>0$ for all $\alpha\in\Delta_{0+}$ or $(\Lambda+\rho|\alpha)\geq 0$ for all $\alpha\in\Delta_{0+}$ and $\mathrm{atp}(\Lambda+\rho)=1$ *then $j_\Lambda=1$.*

Proof. If V is a typical $\mathfrak{g}$-module, then formula (3.3) with $S_\Lambda=\emptyset$ and some $j_\Lambda>0$ follows from the description of singular weights of the Verma module $M(\Lambda)$ and W-anti-invariance of $e^\rho R\,\mathrm{ch}\,V$ (*cf.* [K4]). But if $j_\Lambda\neq 1$, then the stabilizer of $\Lambda+\rho$ in W is non-trivial, and since it is generated by reflections, we would get that $\mathrm{ch}\,V=0$, which is impossible, proving (a).

(c) is clear and (b) follows from [BL, JHKT2, VJ2] (or as above). ■

Theorem 3.1. *Let V be a finite-dimensional irreducible $\mathfrak{g}$-module with highest weight Λ with respect to a subset of positive roots Δ_+. If $S_\Lambda \subset \Pi \subset \Delta_+$, then the $\mathfrak{g}$-module V is tame with respect to (Δ_+, S_Λ).*

Note that Theorem 2.1 follows from Theorem 3.1 and Remark 3.2b.

Example 3.1. If $\operatorname{atp} V = 0$, then $S_\Lambda = \emptyset$ and Theorem 3.1 gives the well-known character formula of a typical $\mathfrak{g}$-module (*cf.* Proposition 3.2a).

Example 3.2. If $\operatorname{atp} V = 1$ and $\mathfrak{g}$ is of the first kind, *i.e.,* of type $A(m,n)$ or $C(n)$, then one can always choose Δ_+ such that $S_\Lambda \subset \Pi$, hence by Theorem 3.1 the $\mathfrak{g}$-module V is tame (*cf.* Proposition 3.2b). There exist, however, modules of atypicality 3 over $A(4,3)$ that are not tame for any choice of Δ_+ and S_Λ (examples were provided by J. Van der Jeugt).

Example 3.3. Let now $\mathfrak{g}$ be of the second kind, *i.e.,* of type $B(m,n)$, $D(m,n)$, $D(2,1;\alpha)$, $G(3)$ or $F(4)$. We exclude $B(0,n)$ for which all V are typical. Let Δ_+ be a standard set of positive roots. Here are the corresponding Dynkin diagrams [K2, p. 56]:


$B(m,n), m > 0$
$\bigcirc - \bigcirc - \ldots - \otimes_2 - \bigcirc_2 - \ldots - \bigcirc_2 \Rightarrow \bigcirc_2$
$B(0,n)$
$\bigcirc - \bigcirc - \ldots - \bigcirc - \bigcirc \Rightarrow \bigcirc$
$D(m,n), m \geq 2$
$\bigcirc - \bigcirc - \ldots - \otimes_2 - \bigcirc_2 - \ldots - \bigcirc_2$
$\bigcirc_1$
$\bigcirc_1$
$D(2,1;\alpha)$
${}^2\otimes$
$\bigcirc_1$
$\bigcirc_1$
$G(3)$
$\otimes_2 - \bigcirc_4 \Lleftarrow \bigcirc_2$
$F(4)$
$\otimes_2 - \bigcirc_3 \Leftarrow \bigcirc_2 - \bigcirc_1$

Let $\tilde{\theta}$ be the only simple root of Δ_{0+} which is not in Π; its coefficients in terms of Π are given above (note that $\tilde{\theta} = \theta$ in all cases except for $B(m,n)$ and $D(m,n)$ with $n > 1$). Let $b = -(\rho|\tilde{\theta}^\vee)$ (as usual, $\alpha^\vee := 2\alpha/(\alpha|\alpha)$ for $\alpha \in \Delta_0$); the values of b are as follows:

$$B(m,n) : m - \tfrac{1}{2}, \quad D(m,n) : m - 1, \quad D(2,1:\alpha) : 1, \quad G(3) : \tfrac{5}{2}, \quad F(4) : 3.$$

Let V be a finite-dimensional irreducible $\mathfrak{g}$-module and let Λ be its highest weight. Let $k = (\Lambda|\tilde{\theta}^\vee)$. Recall that $\dim V < \infty$ implies that $k_i := (\Lambda|\alpha_i^\vee) \in \mathbb{Z}_+$ for all $\alpha_i \in \Pi \bigcap \Delta_0$

and that $k \in \mathbb{Z}_+$. For $k > b$ these conditions are sufficient, and for $k \leq b$ there are some extra conditions listed in [K2, p. 84] (note that b there slightly differs from b here). Suppose that $\operatorname{atp} V = 1$ (this is always the case for a non-typical V over $\mathfrak{g}$ of type $B(m,1)$, $B(1,n)$, $D(m,1)$, $D(2,1;\alpha)$, $G(3)$, and $F(4)$), and let $S_\Lambda = \{\beta\}$ (the choice of β may not be unique). We have: $j_\Lambda = 1$ if $k \geq b$ and $j_\Lambda = 2$ if $k < b$. If the $\mathfrak{g}$-module with the highest weight $r_{\tilde{\theta}}(\Lambda+\rho)-\rho$ is not finite-dimensional (which is always the case when $k \leq b$ or $k > 2b$), then V is tame, *i.e.*,

$$\operatorname{ch} V = \operatorname{ch}_\Lambda .$$

Otherwise we have:

$$\operatorname{ch} V = \operatorname{ch}_\Lambda - \operatorname{ch}_{r_{\tilde{\theta}}(\Lambda+\rho)-\rho} .$$

Furthermore, provided that the highest weight vector is even, we have:

$$\operatorname{sign}(\operatorname{sdim}_\Lambda) = (-1)^{\ell(w)},$$

where $w \in W$ is the shortest element such that $w(\beta) \in \Pi$. Here is a complete list of non-tame modules over exceptional Lie superalgebras (we write $\Lambda = (k; k_2, k_3, \ldots)$):

$$\begin{array}{ll}
D(2,1;\alpha): & \Lambda = \theta = (2;0,0) : \operatorname{ch} V = \operatorname{ch}_\theta - 1. \\
G(3): & \Lambda = (3;0,n) : \operatorname{ch} V = \operatorname{ch}_{(3;0,n)} - \operatorname{ch}_{(2;0,n)} . \\
 & \Lambda = (5;0,0) : \operatorname{ch} V = \operatorname{ch}_{(5;0,0)} - 1. \\
F(4): & \Lambda = (4;0,n,0) : \operatorname{ch} V = \operatorname{ch}_{(4;0,n,0)} - \operatorname{ch}_{(2;0,n,0)} . \\
 & \Lambda = (6;0,0,0) : \operatorname{ch} V = \operatorname{ch}_{(6;0,0,0)} - 1.
\end{array}$$

In the case of $D(2,1;\alpha)$ this result is consistent with [VJ1].

A proof of these results will be given elsewhere. Here we note that the same proof as of Theorem 2.1 shows that for $\mathfrak{g}$ of defect 1 and atypical V with the highest weight Λ such that $(\Lambda+\rho|\beta) = 0$, $\beta \in \bar{\Delta}_{1+}$ and such that $k < b$ for $\mathfrak{g}$ of the second kind one has:

$$e^\rho \mathrm{R} \operatorname{ch} V = \sum_{w \in W^\sharp} \epsilon(w) w \left(e^{\Lambda+\rho}/(1+e^{-\beta})\right) . \tag{3.5}$$

Given integers a and b such that $-1 \leq a \leq b$, we let

$$C(a,b) = \sum_{j=1}^{b} 2^{b-j} \binom{j}{a} \text{ if } a \geq 0; \quad C(-1,b) = \delta_{-1,b} 2^{b+1}. \tag{3.6}$$

Let $N_\epsilon = |\Delta_{\epsilon+}|$, $\epsilon = 0$ or 1.

Theorem 3.2. *Let $\Lambda \in \mathfrak{h}^*$, and let $S_\Lambda = \{\beta_1, \dots, \beta_s\}$. Then (here and further $|\vec{k}|$ stands for $\sum_i k_i$):*

$$\dim_\Lambda = \frac{2^{N_1 - N_0 - s}}{j_\Lambda} \sum_{\substack{\vec{k} \in \mathbb{Z}_+^s \\ |\vec{k}| \leq N_0}} (-1)^{|\vec{k}|} C(|\vec{k}| + s - 1, N_0 + s - 1) \prod_{\alpha \in \Delta_{0+}} \frac{(\Lambda + \rho - \sum_i k_i \beta_i | \alpha)}{(\rho_0 | \alpha)}.$$

The proof of the theorem is based on the following simple lemma.

Lemma 3.1. *Let $P(x_1, \dots, x_s)$ be a polynomial of degree $\leq d$ in the indeterminates $x_1, \dots, x_s$.*

(a) In the domain $|x_i| < 1$ one has

$$\sum_{\vec{n} \in \mathbb{Z}_+^s} (-1)^{|\vec{n}|} \vec{x}^{\vec{n}} P(\vec{n}) = \sum_{\substack{\vec{n} \in \mathbb{Z}_+^s \\ |\vec{n}| \leq d}} \sum_{\substack{\vec{k} \in \mathbb{Z}_+^s \\ k_i \leq n_i}} (-1)^{|\vec{k}|} \binom{\vec{n}}{\vec{k}} P(\vec{k}) \prod_{j=1}^{s} \frac{x_j^{n_j}}{(1 + x_j)^{n_j + 1}}. \tag{3.7}$$

Here and further we use the usual notation:

$$\vec{x}^{\vec{n}} = x_1^{n_1} \dots x_s^{n_s}, \quad \binom{\vec{n}}{\vec{k}} = \binom{n_1}{k_1} \dots \binom{n_s}{k_s}.$$

(b) The value of the rational function on the right-hand side of (3.7) at $x_1 = \ldots = x_s = 1$ equals

$$2^{-s-d} \sum_{\substack{\vec{n} \in \mathbb{Z}_+^s \\ |\vec{n}| \leq d}} (-1)^{|\vec{n}|} C(|\vec{n}| + s - 1, d + s - 1) P(\vec{n}). \tag{3.8}$$

Proof. It suffices to prove (3.7) for a polynomial $P(x) = \binom{x}{j}$, $0 \leq j \leq d$. But this follows from the well-known formulas:

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} \binom{k}{j} = (-1)^n \delta_{n,j},$$

$$\sum_{n \in \mathbb{Z}_+} (-x)^n \binom{n}{j} = \frac{(-x)^j}{(1+x)^{j+1}} \quad \text{in the domain} \quad |x| < 1.$$

(b) follows from (a) and the formula

$$\sum_{\substack{\vec{n} \in \mathbb{Z}_+^s \\ |\vec{n}| = N}} \binom{\vec{n}}{\vec{k}} = \binom{N + s - 1}{|\vec{k}| + s - 1}.$$

■

Remark 3.3. It is clear from the proof that one can take the summation over $\vec{n}$ in the right-hand side of (3.7) and in (3.8) over the set $[P] \in \mathbb{Z}_+^s$ defined as follows. We let $[\vec{x}^{\vec{a}}] = \{\vec{b} \in \mathbb{Z}_+^s | b_i \leq a_i\}$ and let $[P]$ be the union of $[\vec{x}^{\vec{a}}]$ over all the monomials occuring in P with a non-zero coefficient.

Proof of Theorem 3.2. Given $\lambda \in \mathfrak{h}^*$, define the specialization $F_{t\lambda}$ by

$$F_{t\lambda}(e^{\mu}) = e^{t(\lambda|\mu)}, \quad \mu \in \mathfrak{h}^*.$$

First, we compute the following specialization, where $|x_i| < 1$ for all i and $R_0 = \prod_{\alpha \in \Delta_{0+}} (1 - e^{-\alpha})$:

$$\begin{aligned} F_{t\rho_0} & \left(\frac{1}{e^{\rho_0} R_0} \sum_{w \in W} \epsilon(w) w \frac{e^{\Lambda+\rho}}{\prod_i (1 + x_i e^{-\beta_i})} \right) \\ &= \sum_{\vec{n} \in \mathbb{Z}_+^s} (-1)^{|\vec{n}|} \vec{x}^{\vec{n}} \frac{\displaystyle\sum_{w \in W} \epsilon(w) e^{t(w(\Lambda+\rho-n_1\beta_1-\ldots-n_s\beta_s)|\rho_0)}}{F_{t\rho_0}(e^{\rho_0} R_0)} \\ &= \sum_{\vec{n} \in \mathbb{Z}_+^s} (-1)^{|\vec{n}|} \vec{x}^{\vec{n}} \frac{F_{t(\Lambda+\rho-\Sigma n_i\beta_i)}(e^{\rho_0} R_0)}{F_{t\rho_0}(e^{\rho_0} R_0)}. \end{aligned}$$

Hence we have in the domain $|x_i| < 1$:

$$\begin{aligned} & \lim_{t \to 0} F_{t\rho_0} \left(\frac{1}{e^{\rho_0} R_0} \sum_{w \in W} \epsilon(w) w \frac{e^{\Lambda+\rho}}{\prod_i (1 + x_i e^{-\beta_i})} \right) \\ & = \sum_{\vec{n} \in \mathbb{Z}_+^s} (-1)^{|\vec{n}|} \vec{x}^{\vec{n}} \prod_{\alpha \in \Delta_{0+}} \frac{(\Lambda + \rho - \sum_i n_i\beta_i | \alpha)}{(\rho_0|\alpha)}. \end{aligned} \tag{3.9}$$

Note that the expression under the limit at the left-hand side of (3.9) is a continuous function in the domain $|t| < \epsilon$ and $|x_i - 1| < \epsilon$ for sufficiently small ϵ. Hence we may calculate

$$\lim_{t \to 0} F_{t\rho_0} \left(\frac{1}{e^{\rho_0} R_0} \sum_{w \in W} \epsilon(w) w \frac{e^{\Lambda+\rho}}{\prod_i (1 + e^{-\beta_i})} \right)$$

by applying Lemma 3.1a to the right-hand side of (3.9) and then evaluating it at $x_1 = \ldots = x_s = 1$ using Lemma 3.2b. Since $\dim_\Lambda = \lim_{t \to 0} F_{t\rho_0} \operatorname{ch}_\Lambda$, we obtain the result. ■

Theorem 3.3. *Let $\Lambda \in \mathfrak{h}^*$, let S_Λ be a $\Lambda+\rho$-isotropic subset and let $s = |S_\Lambda|$. Consider the root system $M_\Lambda := \{\alpha \in \bar{\Delta}_0 | \alpha \perp S_\Lambda\}$, let ρ^0_Λ be the half of the sum of roots from $M^+_\Lambda := M_\Lambda \bigcap \Delta_+$ and let $L_\Lambda(\lambda)$ denote the irreducible module with highest weight λ over* *the reductive Lie algebra with root system* M_Λ. *Let* $a = n - m - 1$ *for* $\mathfrak{g}$ *of type* $B(m,n)$ *with* $m < n$ *and let* $a = 0$ *in all other cases. Then*

$$|\mathrm{sdim}_\Lambda| = |W/W^\sharp| 2^{-a} j_\Lambda^{-1} \dim L_\Lambda(\Lambda + \rho - \rho_\Lambda^0) \text{ if } s = \operatorname{def}\mathfrak{g}, \tag{3.10a}$$

$$\mathrm{sdim}_\Lambda = 0 \text{ otherwise.} \tag{3.10b}$$

Proof. Let W^Λ be the subgroup of W generated by reflections with respect to the roots from M_Λ and let W_1 be a set of left coset representatives, so that $W = W_1 W^\Lambda$. Due to (2.3), formula (3.4) can be rewritten using the Weyl character formula for the root system M_Λ as follows:

$$j_\Lambda \prod_{\alpha\in\Delta_{1+}\setminus\bar{\Delta}_{1+}} (1+e^{-\alpha}) \prod_{\alpha\in\bar{\Delta}_{0+}} (1-e^{-\alpha}) e^\rho \,\mathrm{sch}_\Lambda$$
$$= \prod_{\alpha\in\bar{\Delta}_{1+}} (1-e^{-\alpha}) \sum_{w\in W_1} \check{\epsilon}(w) \frac{\prod_{\alpha\in M_\Lambda^+}(1-e^{-w(\alpha)})}{\prod_{\alpha\in S_\Lambda}(1-e^{-w(\alpha)})} w\left(e^{\rho_\Lambda^0} \operatorname{ch} L_0(\Lambda+\rho-\rho_\Lambda^0)\right).$$

For $\lambda \in \mathfrak{h}^*$ such that $(\lambda|\alpha) \neq 0$ for all $\alpha \in \Delta$, we apply to both sides of this equality the specialization $F_{t\lambda}$ to obtain as $t \to 0$:

$$j_\Lambda 2^{|\Delta_{1+}\setminus\bar{\Delta}_{1+}|} \prod_{\alpha\in\bar{\Delta}_{0+}\setminus\bar{\Delta}_{1+}} (\lambda|\alpha)\,\mathrm{sdim}_\Lambda =$$
$$t^{|\bar{\Delta}_{1+}|+|M_\Lambda^+|-|\bar{\Delta}_{0+}|-|S_\Lambda|} \left(\sum_{w\in W_1} \check{\epsilon}(w) \prod_{\alpha\in M_\Lambda^+\setminus S_\Lambda} (\lambda|w(\alpha))\right) \dim L_\Lambda(\Lambda+\rho-\rho_\Lambda^0) + o(t).$$

Due to Lemma 2.2d, if $s < \operatorname{def}\mathfrak{g}$, the right-hand side tends to 0 as $t \to 0$, proving (3.10b), and if $s = \operatorname{def}\mathfrak{g}$, we obtain as $t \to 0$:

$$\mathrm{sdim}_\Lambda = j_\Lambda^{-1} c(\lambda, \Delta_+, S_\Lambda) \dim L_\Lambda(\Lambda+\rho-\rho_\Lambda^0), \tag{3.11}$$

where

$$c(\lambda,\Delta_+,S_\Lambda) = \prod_{\alpha\in\bar{\Delta}_{0+}\setminus\bar{\Delta}_{1+}} (\lambda|\alpha) \sum_{w\in W_1} \check{\epsilon}(w) \prod_{\alpha\in M_\Lambda^+\setminus S_\Lambda} (\lambda|w(\alpha)) \tag{3.12}$$

is a constant which, in fact, is independent of the choice of λ (since all other factors in (3.12) are independent of λ). It follows that the constant $c = c(\lambda, \Delta_+, S_\Lambda)$ is independent up to a sign of the choice of (Δ_+, S_Λ) as well. Recall now that, according to Theorem 2.1, the 1-dimensional module is tame with respect to a certain pair (Δ'_+, S'). Applying (3.11) to the 1-dimensional $\mathfrak{g}$-module (for which $\operatorname{sdim} = \pm 1$) and using (3.4), we obtain:

$$1 = \pm |W/W^\sharp|^{-1} c \dim L_0(\rho' - \rho_0^{0\prime}).$$

Finally a case by case inspection shows that $\dim L_0(\rho' - \rho_0^{0\prime}) = 2^a$, proving (3.10a).

In the case $\mathfrak{g} = C(n)$ a formula equivalent to (3.10b) was obtained in [VJ2]. ■

Conjecture 3.1. $\operatorname{sdim} V = 0$ if and only if $\operatorname{atp} V < \operatorname{def} \mathfrak{g}$.

Example 3.4. (a) if V is a typical $\mathfrak{g}$-module then formulas (3.6) and (3.10) turn into the known formulas [K4]:

$$\dim V = 2^{N_1} \prod_{\alpha \in \Delta_{0+}} \frac{(\Lambda + \rho|\alpha)}{(\rho_0|\alpha)},$$

$$\operatorname{sdim} V = 0 \text{ if } \mathfrak{g} \neq B(0,n), = \prod_{\alpha \in \bar{\Delta}_{0+}} (\Lambda + \rho|\alpha)/(\rho|\alpha) \text{ if } \mathfrak{g} = B(0,n).$$

(b) Let $\mathfrak{g}$ be an exceptional Lie superalgebra and let V be a finite-dimensional irreducible non-typical $\mathfrak{g}$-module with highest weight Λ, so that $\operatorname{atp} V = 1$. We keep the notation of Example 3.3.

$\mathfrak{g} = D(2,1;\alpha) : (\alpha_1|\alpha_1) = 0$, $(\alpha_1|\alpha_2) = -1$, $(\alpha_1|\alpha_3) = -\alpha$, $(\alpha_2|\alpha_2) = 2$, $(\alpha_3|\alpha_3) = 2\alpha$, $(\alpha_2|\alpha_3) = 0$. Assume that V is neither the 1-dimensional nor the adjoint module. Then $\operatorname{sdim} V = \pm 2$ and $\frac{1}{2} \dim V$ is given by the following formulas, where $m = k_2$, $n = k_3$:

$$4kmn + 2k(m+n) - 2n + 1 \text{ if } \beta = \alpha_2,$$

$$4kmn + 2(km + 3kn - mn) + 4(k-n) - 1 \text{ if } \beta = \alpha_1 + \alpha_2,$$

$$4kmn + 2(3km + kn - mn) + 4(k-m) - 1 \text{ if } \beta = \alpha_2 + \alpha_3,$$

$$4kmn + 2(3km + 3kn - mn) + 4(2k - m - n) - 7 \text{ if } \beta = \alpha_1 + \alpha_2 + \alpha_3.$$

(Recall that for a typical V one has [K4]: $\dim V = 16(m+1)(n+1)(k-1)$ and $\operatorname{sdim} V = 0$.)

$\mathfrak{g} = G(3)$ $(m = k_2,\ n = k_3)$:

If $\beta = \alpha_1$, then $\dim V = 1$; $\beta \neq \alpha_1 + \alpha_2$. If $\beta = \alpha_1 + \alpha_2 + \alpha_3$, then $k = 2$, $m = 0$ and $\operatorname{sdim} V = 2n+3$. If $\beta = \alpha_1 + 3\alpha_2 + \alpha_3$, then $2k = m+6$; for $k = 3$ we have $\operatorname{sdim} V = -2n-3$, for $k>3$ we have $\operatorname{sdim} V=-2(m+2n+3)$. If $\beta=\alpha_1+3\alpha_2+2\alpha_3$, then $2k=m+3n+9$ and $\operatorname{sdim} V=2(m+n+2)$. If $\beta=\alpha_1+4\alpha_2+2\alpha_3$, then $2k=2m+3n+10$; for $k=5$ we have $\dim V=321$, $\operatorname{sdim} V=-1$, for $k>5$ we have $\operatorname{sdim} V=-2(n+1)$.

$\mathfrak{g}=F(4)$ ($l=k_2$, $m=k_3$, $n=k_4$):

If $\beta=\alpha_1$, then $\dim V=1$; $\beta\neq\alpha_1+\alpha_2$. If $\beta=\alpha_1+\alpha_2+\alpha_3$, then $k=2$, $l=n=0$ and $\operatorname{sdim} V=(m+2)^3$. If $\beta=\alpha_1+2\alpha_2+\alpha_3$, then $3k=l-2n+8$; if $\beta=\alpha_1+\alpha_2+\alpha_3+\alpha_4$, then $3k=2n-l+10$; in both cases $k\geq 3$ and $\operatorname{sdim} V=-(m+1)(l+m+n+3)(l+2m+n+4)$. If $\beta=\alpha_1+2\alpha_2+\alpha_3+\alpha_4$, then $3k=l+2n+12$; if $k=4$, then $l=n=0$ and $\operatorname{sdim} V=(m+2)^3$; if $k>4$, then $\operatorname{sdim} V=(l+m+2)(m+n+2)(l+2m+n+4)$. If $\beta=\alpha_1+2\alpha_2+2\alpha_3+\alpha_4$, then $3k=l+4m+2n+16$ and $\operatorname{sdim} V=-(l+m+2)(n+1)(l+m+n+3)$. If $\beta=\alpha_1+3\alpha_2+2\alpha_3+\alpha_4$, then $3k=3l+4m+2n+18$; if $k=6$, then $l=m=n=0$ and $\operatorname{sdim} V=1$; if $k>6$, then $\operatorname{sdim} V=(m+1)(n+1)(m+n+2)$.

§4. The denominator identity for affine superalgebras and its specializations.

Let $\mathfrak{g}$, as before, be a finite-dimensional simple Lie superalgebra over $\mathbb{C}$ of rank ℓ with a non-degenerate even invariant bilinear form $(.|.)$ (see Section 1). Let $\hat{\mathfrak{g}}=\mathbb{C}[t,t^{-1}]\otimes\mathfrak{g}\oplus\mathbb{C}C\oplus\mathbb{C}d$ be the associated to $\mathfrak{g}$ affine superalgebra. The $\mathbb{Z}_2$-gradation of $\hat{\mathfrak{g}}$ extends from that of $\mathfrak{g}$ by letting $\deg t=0$, $\deg C=\deg d=0$. Denoting $a(n)=t^n\otimes a$ for $a\in\mathfrak{g}$, $n\in\mathbb{Z}$, we have the following commutation relations ($a,b\in\mathfrak{g}$; $m,n\in\mathbb{Z}$):

$$[a(m),b(n)]=[a,b](m+n)+m\delta_{m,-n}(a|b)C,$$
$$[d,a(m)]=ma(m),\quad [C,\hat{\mathfrak{g}}]=0.$$

We identify $\mathfrak{g}$ with the subalgebra $1\otimes\mathfrak{g}$ of $\hat{\mathfrak{g}}$. The bilinear form $(.|.)$ extends from $\mathfrak{g}$ to an even invariant bilinear form on $\hat{\mathfrak{g}}$ by:

$$(a(m)|b(n))=\delta_{m,-n}(a|b),\qquad \big(\mathbb{C}[t,t^{-1}]\otimes\mathfrak{g}|\mathbb{C}C+\mathbb{C}d\big)=0,$$
$$(C|C)=(d|d)=0,\qquad (C|d)=1.$$

Let $\hat{\mathfrak{h}}=\mathfrak{h}+\mathbb{C}C+\mathbb{C}d$ be the Cartan subalgebra of $\hat{\mathfrak{g}}$ and let $\hat{\mathfrak{n}}_+=\mathfrak{n}_+\oplus\Big(\bigoplus_{j>0}t^j\otimes\mathfrak{h}\Big)$. We identify $\hat{\mathfrak{h}}^*$ with $\hat{\mathfrak{h}}$ using the bilinear form $(.|.)$.

Given a subset $S \subset \mathfrak{h}$, we let $\tilde{S} = \{\alpha + nC | \alpha \in S, n \in \mathbb{Z}\} \setminus \{0\}$. Then the set of roots of $\hat{\mathfrak{g}}$ with respect to $\hat{\mathfrak{h}}$ is $\hat{\Delta} = \tilde{\Delta}_0 \cup \{0\}\tilde{}.$, the set of even (resp. odd) roots being $\hat{\Delta}_0 = \tilde{\Delta}_0 \cup \{0\}\tilde{}$ (resp. $\hat{\Delta}_1 = \tilde{\Delta}_1$). The set of positive roots of $\hat{\Delta}$ associated to the choice of $\Delta_+ \subset \Delta$ is $\hat{\Delta}_+ = \Delta_+ \cup \{\alpha + nC | \alpha \in \Delta \cup \{0\}, n > 0\}$, the set of simple roots being $\hat{\Pi} = \{\alpha_0 := C - \theta\} \cup \Pi$. As usual the subscript $+$ means the intersection with $\hat{\Delta}_+$, *e.g.*, $\tilde{\Delta}_+ = \tilde{\Delta} \cap \hat{\Delta}_+$.

Define $\hat{\rho} \in \hat{\mathfrak{h}}^*$ by $\hat{\rho} = \rho + h^\vee d$. One has:

$$(\hat{\rho}|\alpha_i) = \tfrac{1}{2}(\alpha_i|\alpha_i) \text{ for } \alpha_i \in \hat{\Pi}, \quad (\hat{\rho}|d) = 0, \quad (\hat{\rho}|C) = h^\vee.$$

As before, for $\alpha \in \tilde{\Delta}_0$ we let $\alpha^\vee = 2\alpha/(\alpha|\alpha)$. For $\alpha \in \mathfrak{h}^*$ define $t_\alpha \in \operatorname{Aut} \hat{\mathfrak{h}}^*$ by

$$t_\alpha(\lambda) = \lambda + \lambda(C)\alpha - \big((\lambda|\alpha) + \tfrac{1}{2}(\alpha|\alpha)\lambda(C)\big)\, C.$$

For an additive subgroup $S \subset \mathfrak{h}^*$ the set $t_S := \{t_\alpha | \alpha \in S\}$ is a subgroup of $\operatorname{Aut} \hat{\mathfrak{h}}^*$. Let M(resp$M^\sharp$) be the $\mathbb{Z}$-span of the $\alpha^\vee$ such that $\alpha \in \Delta_0$ (resp. $\alpha \in \Delta_0^\sharp$). The Weyl group of $\hat{\mathfrak{g}}$ is $\widehat{W} = W \ltimes t_M$. The following subgroup of $\widehat{W}$ is more important, however: $\widehat{W}^\sharp = W \ltimes t_{M^\sharp}$.

Denote by $\widehat{\mathcal{F}}$ the field of meromorphic functions in the domain $Y = \{h \in \hat{\mathfrak{h}} | \operatorname{Re}(C|h) > 0\}$. Since C is $\widehat{W}$-fixed, the group $\widehat{W}$ acts on $\widehat{\mathcal{F}}$. Let $q = e^{-C}$; note that $|q| < 1$ in Y. Given $\lambda \in \mathfrak{h}^*$, we extend it to a linear function $\tilde{\lambda} \in \hat{\mathfrak{h}}^*$ by letting $\tilde{\lambda}(C) = \tilde{\lambda}(d) = 0$; this gives us an embedding of the field $\mathcal{F}$ of rational exponential functions on $\mathfrak{h}$ in the field $\widehat{\mathcal{F}}$.

Let R be the Weyl denominator of $\mathfrak{g}$ (see Section 2). Introduce the *affine denominator* by

$$\widehat{\mathrm{R}} = \mathrm{R} \prod_{n=1}^{\infty} (1 - q^n)^\ell \prod_{\alpha \in \Delta_0} (1 - q^n e^\alpha) \prod_{\alpha \in \Delta_1} (1 + q^n e^\alpha)^{-1}.$$

It is clear that $\widehat{\mathrm{R}} \in \widehat{\mathcal{F}}$. As in Section 2, it is easy to check that

$$w(e^{\hat{\rho}}\widehat{\mathrm{R}}) = \epsilon(w) e^{\hat{\rho}}\widehat{\mathrm{R}}, \quad w \in \widehat{W}. \tag{4.1}$$

Here $\epsilon : \widehat{W} \to \pm 1$ is a homomorphism which extends the homomorphism $\epsilon : W \to \pm 1$ by $\epsilon(t_M) = 1$.

Theorem 4.1. *Suppose that* $h^\vee \neq 0$. *Then*

$$e^{\hat\rho}\widehat{\mathrm{R}} = \sum_{\alpha\in M^\sharp} t_\alpha(e^{\hat\rho}\mathrm{R}). \tag{4.2}$$

The proof of Theorem 4.1 is based on the following

Lemma 4.1. *Suppose that* $\Lambda\in\hat{\mathfrak{h}}^*$ *is such that* $(\Lambda+\hat\rho)(C)\neq 0$, *and suppose that* $\hat L(\lambda)$, $\lambda\in\hat{\mathfrak{h}}^*$, *is an irreducible subquotient of the Verma* $\hat{\mathfrak{g}}$*-module* $\widehat{\mathrm{M}}(\Lambda)$. *Then there exists a sequence* $\lambda_0=\Lambda,\lambda_1,\dots,\lambda_k=\lambda$ *such that* λ_i *is obtained from* λ_{i-1} $(0<i\leq k)$ *in one of the following two ways:*

(i) $\lambda_i=\lambda_{i-1}-n\beta_i$, *where* $\beta_i\in\tilde\Delta_{0+}$ *and* $(\lambda_{i-1}+\hat\rho|\beta_i^\vee)=n\in\mathbb{N}$,

(ii) $\lambda_i=\lambda_{i-1}-\beta_i$, *where* $\beta_i\in\tilde\Delta_{1+}$, $(\beta_i|\beta_i)=0$, *and* $(\lambda_{i-1}+\hat\rho|\beta_i)=0$.

Proof. See [K6, Theorem 1b]. (Note that the usual proof is based on the determinant formula given by [K6, Theorem 1a]; the exponents in this formula should be corrected as in [K8]). ■

A detailed proof of Theorem 4.1 will be given elsewhere. In the case def $\mathfrak{g}=1$ the same proof works as that in Section 2.

Theorem 4.2. *Let* $\mathfrak{g}$ *be a simple finite-dimensional Lie superalgebra with a non-degenerate Killing form,* i.e., $h^\vee\neq 0$. *Choose a set of positive roots* Δ_+ *such that the set of simple roots contains a maximal set of pairwise orthogonal isotropic roots* $\{\beta_1,\dots,\beta_s\}$ $(s=\operatorname{def}\mathfrak{g})$. *Let* $\beta=\sum_{i=1}^s\beta_i$ *and let* $d_\epsilon=\dim\mathfrak{g}_\epsilon$, $\epsilon=0$ *or* 1. *Then*

$$\prod_{j=1}^\infty\frac{(1-q^j)^{d_0}}{(1+q^j)^{d_1}} = \frac{2^{N_1}}{|W/W^\sharp|}\sum_{\alpha\in M^\sharp} q^{\frac{1}{2}h^\vee(\alpha|\alpha)+(\rho+\beta|\alpha)} \sum_{\substack{\vec n\in\mathbb{Z}_+^s\\ |\vec n|\leq N_0}} \frac{A_\alpha(\vec n)}{\prod_i(1+q^{(\alpha|\beta_i)})^{n_i+1}}, \tag{4.3}$$

where

$$A_\alpha(\vec n)=\sum_{\substack{\vec k\in\mathbb{Z}_+^s\\ k_i\leq n_i}}(-1)^{|\vec k|}\binom{\vec n}{\vec k}\prod_{\gamma\in\Delta_{0+}}\frac{(\rho+h^\vee\alpha-k_1\beta_1-\dots-k_s\beta_s|\gamma)}{(\rho_0|\gamma)}. \tag{4.4}$$

Proof. From Theorems 2.1 and 4.1, and Remark 3.2c we have:

$$\frac{|W/W^\sharp|\widehat{\mathrm{R}}}{e^{\rho_0}\mathrm{R}_0} = \lim_{x_1,\dots,x_s\to 1}\frac{e^{-\hat\rho}}{e^{\rho_0}\mathrm{R}_0}\sum_{\alpha\in M^\sharp}\sum_{w\in W}\epsilon(w)w\,t_\alpha\frac{e^{\hat\rho}}{\prod_j(1+x_je^{-\beta_j})}. \tag{4.5}$$

Assuming that $|x_j|$ are sufficiently small we expand in a geometric progression for each α:

$$t_\alpha \frac{e^{\hat\rho}}{\prod_j (1+x_j e^{-\beta_j})} = q^{\frac{1}{2}h^\vee(\alpha|\alpha)+(\rho|\alpha)} \sum_{\vec n\in\mathbb{Z}_+^s} \prod_j \left(-x_j q^{-(\alpha|\beta_j)}\right)^{n_j} e^{\hat\rho+h^\vee\alpha-\sum_j n_j\beta_j}.$$

Hence we may rewrite (4.5) as follows:

$$\begin{aligned}\frac{|W/W^\sharp|\widehat{\mathrm{R}}}{e^{\rho_0}\mathrm{R}_0} &= \lim_{x_1,\dots,x_s\to 1}\sum_{\alpha\in M^\sharp} q^{\frac{1}{2}h^\vee(\alpha|\alpha)+(\rho|\alpha)} \\ &\qquad\times\sum_{\vec n\in\mathbb{Z}_+^s}\prod_j (-x_j q^{-(\alpha|\beta_j)})^{n_j} \frac{\sum_{w\in W}\epsilon(w)e^{w(\hat\rho+h^\vee\alpha-\sum_j n_j\beta_j)-\hat\rho}}{e^{\rho_0}\mathrm{R}_0}.\end{aligned} \tag{4.6}$$

Applying the specialization $F_{t\rho_0}$ to both sides and tending t to 0 we obtain:

$$\begin{aligned}&\frac{|W/W^\sharp|}{2^{N_1}}\prod_{n=1}^\infty \frac{(1-q^n)^{\dim\mathfrak{g}_0}}{(1+q^n)^{\dim\mathfrak{g}_1}} \\ &= \lim_{x_1,\dots,x_s\to 1}\sum_{\alpha\in M^\sharp} q^{\frac{1}{2}h^\vee(\alpha|\alpha)+(\rho|\alpha)} \sum_{\vec n\in\mathbb{Z}_+^s}\prod_j\left(-x_j q^{-(\alpha|\beta_j)}\right)^{n_j}\prod_{\gamma\in\Delta_{0+}}\frac{(\rho+h^\vee\alpha-\sum_j n_j\beta_j|\gamma)}{(\rho_0|\gamma)}.\end{aligned}$$

The theorem now follows from (3.7). ∎

Theorem 4.3. *Let $\mathfrak{g}$ be a simple finite-dimensional Lie superalgebra of type $A(m,n)$, $m\neq n$, or $C(m)$. Then in the notation of Theorem 4.2 one has:*

$$\prod_{j=1}^\infty \frac{(1-q^j)^{d_0}}{(1+q^{j-\frac{1}{2}})^{d_1}} = \frac{q^{r+\frac{1}{2}s}}{|W/W^\sharp|}\sum_{\alpha\in M^\sharp} q^{\frac{1}{2}h^\vee(\alpha|\alpha)+(\rho+\beta|\alpha)} \sum_{\substack{\vec n\in\mathbb{Z}_+^s\\ |\vec n|\le N_0}} \frac{A_\alpha(\vec n)}{\prod_i\left(1+q^{(\alpha|\beta_i)+\frac{1}{2}}\right)^{n_i+1}}, \tag{4.7}$$

where r is the number of roots from Δ_{1+} which are negative with respect to the standard set of positive roots.

Proof. Let $\lambda\in\mathfrak{h}^*$ be such that $(\lambda|\alpha)=0$ if $\alpha\in\Delta_0$ and $(\lambda|\alpha)=\frac{1}{2}$ if α is odd and positive with respect to the standard set of positive roots. Now apply to both sides of (4.6) the specialization $F_{t\rho_0-(\log q)\lambda}$ and tend t to 0. ∎

Example 4.1. Let $\mathfrak{g} = sl(2,1)$. Choose Δ_+ such that both simple roots α_1 and α_2 are odd. Then $(\alpha_1|\alpha_1) = (\alpha_2|\alpha_2) = 0$, $(\alpha_1|\alpha_2) = 1$, $\rho = 0$, $h^\vee = 1$, $M^\sharp = \mathbb{Z}(\alpha_1 + \alpha_2)$. We let $x = e^{-\alpha_1}$, $y = e^{-\alpha_2}$. Then (2.4) becomes (if we take $\beta = \alpha_1$ or α_2 respectively):

$$R := \frac{1-xy}{(1+x)(1+y)} = \frac{1}{1+x} - \frac{y}{1+y} = \frac{1}{1+y} - \frac{x}{1+x}.$$

Formula (4.2) for $\widehat{\mathrm{R}}$ becomes after expanding the right-hand side (for which we assume $|q| < |x|$, $|y| < 1$):

$$\begin{aligned} &\prod_{n=1}^{\infty} \frac{(1-q^n)^2(1-xyq^{n-1})(1-x^{-1}y^{-1}q^n)}{(1+xq^{n-1})(1+x^{-1}q^n)(1+yq^{n-1})(1+y^{-1}q^n)} \\ &= \left(\sum_{m,n=0}^{\infty} - \sum_{m,n=-1}^{-\infty} \right) \left((-1)^{m+n} x^m y^n q^{mn} \right). \end{aligned} \tag{4.8}$$

This truly remarkable identity may be viewed also as the denominator identity for $N = 4$ superconformal algebra. An equivalent form of (4.8) is (here we assume $|q| < |x| < 1$):

$$\prod_{k=1}^{\infty} \frac{(1-q^n)^2(1-xyq^{n-1})(1-x^{-1}y^{-1}q^n)}{(1+xq^{n-1})(1+x^{-1}q^n)(1+yq^{n-1})(1+y^{-1}q^n)} = \sum_{j\in\mathbb{Z}} \frac{(-x)^j}{1+yq^j}.$$

Replacing in this formula q by q^2, x by z and y by qz^{-1}, we obtain

$$\prod_{n=1}^{\infty} \frac{(1-q^n)^2}{(1+q^{n-1}z)(1+q^n z^{-1})} = \sum_{n\in\mathbb{Z}} (-1)^n \frac{q^{n(n+1)/2}}{1+q^n z}.$$

This identity may be viewed as the denominator identity for $N = 2$ superconformal algebra and also of the affine superalgebra $gl(1,1)^\wedge$. It also has been rediscovered many times (see, *e.g.,* [KP]). It is worth mentioning here that Gauss identities (5.1) and (5.2) are denominator identities for $N = 1$ superconformal algebras.

Yet another useful form of (4.8) is obtained by dividing both sides by R:

$$\begin{aligned} &\prod_{n=1}^{\infty} \frac{(1-q^n)^2(1-xyq^n)(1-x^{-1}y^{-1}q^n)}{(1+xq^n)(1+x^{-1}q^n)(1+yq^n)(1+y^{-1}q^n)} \\ &= 1 + \frac{(1+x)(1+y)}{(1-xy)} \sum_{m,n=1}^{\infty} (-1)^{m+n} q^{mn} (x^m y^n - x^{-m} y^{-n}). \end{aligned} \tag{4.9}$$

Letting $x = y = z$, we obtain:

$$\begin{aligned}(4.10)\qquad &\prod_{n=1}^{\infty} \frac{(1-q^n)^2(1-z^2q^n)(1-z^{-2}q^n)}{(1+zq^n)^2(1+z^{-1}q^n)^2} \\ &\quad = 1 + (1+z)\sum_{m,n=1}^{\infty} (-1)^{m+n} q^{mn} \frac{z^{m+n} - z^{-m-n}}{1-z}.\end{aligned}$$

Example 4.2. Let $\mathfrak{g} = \mathrm{osp}(3,2)$. Choose Δ_+ such that both simple roots α_1 and α_2 are odd; one has: $(\alpha_1|\alpha_2) = 0$, $(\alpha_2|\alpha_2) = \frac{1}{2}$, $(\alpha_1|\alpha_2) = -\frac{1}{2}$, $\rho = -\frac{1}{2}\alpha_1$, $h^\vee = \frac{1}{2}$, $M^\sharp = 2\mathbb{Z}\alpha_2$. We let $x = e^{-\alpha_1}$, $y = e^{-\alpha_2}$. Then (2.4) becomes:

$$\mathrm{R} := \frac{(1-y)(1-xy)}{(1+x)(1+xy^2)} = \frac{1}{1+x} - \frac{y}{1+xy^2}.$$

Formula (4.2) for $\widehat{\mathrm{R}}$ becomes after expanding the right-hand side (as before, $|q| < |x|$, $|y| < 1$):

$$\begin{aligned}&\prod_{n=1}^{\infty} \frac{(1-q^n)^2(1-xyq^{n-1})(1-x^{-1}y^{-1}q^n)(1-y^2q^{n-1})(1-y^{-2}q^n)}{(1+xq^{n-1})(1+x^{-1}q^n)(1+yq^{n-1})(1+y^{-1}q^n)(1+xy^2q^{n-1})(1+x^{-1}y^{-2}q^n)} \\ (4.11)\qquad &= \left(\sum_{m,n=0}^{\infty} - \sum_{m,n=-1}^{-\infty}\right) (-x)^n \left(y^{-m} - y^{m+2n+1}\right) q^{\frac{1}{2}m(m+2n+1)}.\end{aligned}$$

Here is a general construction of a specialization (*cf.* [K9]). Let σ be an automorphism of order m of $\mathfrak{g}$ such that $\sigma(h) = h$ for $h \in \mathfrak{h}$, let $\mathfrak{g} = \bigoplus_{j \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}_{\epsilon,j}$ ($\epsilon = 0$ or 1) be the corresponding $\mathbb{Z}/m\mathbb{Z}$-gradations and let $d_{\epsilon,j} = \dim \mathfrak{g}_{\epsilon,j}$. For $\alpha \in \Delta$ we have $\mathfrak{g}_\alpha \subset \mathfrak{g}_{\epsilon,\bar{s}_\alpha}$; let $s_\alpha \in \mathbb{Z}$ be such that $0 \leq s_\alpha < m$, $\bar{s}_\alpha = s_\alpha \bmod m$. Define $\lambda \in \mathfrak{h}$ by $(\lambda|\alpha) = s_\alpha$. Replace q by q^m in (4.2), divide both sides of it by $\prod_{\substack{\alpha \in \Delta_{0+} \\ (\lambda|\alpha)=0}} (1 - e^{-\alpha})$ and apply to the resulting identity the specialization $F^\sigma := F_{-(\log q)\lambda}$. The left-hand side becomes

$$\prod_{j \in \mathbb{N}} (1-q^j)^{d_{0,j}} / (1+q^j)^{d_{1,j}}.$$

One can generalize the trick applied in the proof of Theorems 4.2 and 4.3 to evaluate the right-hand side (as in [K9]). However, the answer is simpler when λ is fixed under the action of W (as in Theorems 4.2 and 4.3). Another situation when this trick may be applied is when $s_\alpha = 0$ for $\alpha \in \Delta_0^\sharp$ and $s_\alpha \neq 0$ for $\alpha \in \Delta_0 \setminus \Delta_0^\sharp$, so that λ is fixed under $W^\sharp$.

Example 4.3. (a) Let σ be the automorphism of $\mathfrak{g}$ such that $\sigma|_{\mathfrak{g}_0} = 1$, $\sigma|_{\mathfrak{g}_1} = -1$. Then applying F^σ to $\widehat{\mathrm{R}}/\mathrm{R}_0^\sharp$ gives an expansion of $\prod_{j\in\mathbb{N}}(1-q^{2j})^{d_0}/(1-q^{2j-1})^{d_1}$, which is $\Delta(q)^{d_0}$ (see (5.2) below) provided that $d_0 = d_1$.

(b) Let $\mathfrak{g} = A(n, n-1)$ and choose Δ_+ such that Π consists of odd roots $\{\alpha_1, \ldots, \alpha_{2n}\}$ (see Example 1.1). Let σ be such that $s_{\alpha_{2j}} = 1$ and $s_{\alpha_{2j-1}} = 0$ ($j = 1, \ldots, n$). Then F^σ gives an expansion of $\square(q)^{2n}$ over an n-dimensional lattice. Namely, using the same method as that in the proof of Theorem 4.2, we obtain:

$$\square(-q)^{2n} = 2^n \sum_{\alpha\in M^\sharp} \sum_{s\in\mathbb{S}} (-1)^{|s|} q^{\frac{n+1}{2}(\alpha|\alpha)+(\alpha|\rho_0)-(\alpha|\tilde{s})} \prod_{j=1}^{n} \left(1 + q^{(n+1)(\alpha|\beta_j)-(\tilde{s}|\beta_j)}\right)^{-1},$$

where $\mathbb{S}$ is the set of all subsets in Δ_{0+}, and $|s|$ (resp. $\tilde{s}$) stands for the number (resp. sum) of elements of $s \in \mathbb{S}$. Note that $M^\sharp$ is an n-dimensional lattice (of type A_n), so that for $n = 1$ we recover Jacobi's formula (0.1), but for $n = 2$, 3 and 4 our formula is more complicated than Jacobi's (0.2–4). The way to recover Jacobi's formulas will be explained below.

(c) Let $\mathfrak{g} = A(2n+1, 2n) = sl(2n+2, 2n+1)$ and let σ be the conjugation by the matrix $\mathrm{diag}(I_{n+1}, -I_{n+1};\ I_{n+1}, -I_n)$. Then σ is an order 2 automorphism of $\mathfrak{g}$ such that $d_{0,0} = d_{0,1} = d_{1,0} = d_{1,1} = 2(n+1)(2n+1)$. Hence F^σ gives an expansion of $\square(q)^{2(n+1)(2n+1)}$ over an $n+1$-dimensional lattice ($n = 0, 1, \ldots$).

Example 4.4. Let $\mathfrak{g} = A(m, n)$ with $m > n$. We choose the set of simple roots as in Example 1.1. Then formula (4.2) can be written, using (2.4) and (3.4), as follows:

$$e^{\hat\rho}\hat{\mathrm{R}} = \frac{1}{(n+1)!} \sum_{\alpha\in M^\sharp} \sum_{w\in W} \epsilon(w) w t_{-\alpha} \left(e^{\hat\rho} \prod_{j=1}^{n+1} \left(1 + e^{-\alpha_{2j-1}}\right)^{-1} \right), \tag{4.12}$$

where $M^\sharp$ is the lattice spanned over $\mathbb{Z}$ by elements $\alpha_{2j-1} + \alpha_{2j}$ ($j = 1, \ldots, n+1$) and α_i ($2n+3 \le i \le m+n+1$). Introduce the element $\xi \in \mathfrak{h}$ by relations $(\xi|\alpha_j) = (-1)^j/2$ for $1 \le j \le 2(n+1)$, $(\xi|\alpha_j) = 0$ for $2n+3 \le j \le m+n+1$. Then $(\xi|\alpha) = 0$ for all $\alpha \in \Delta_0$, hence t_ξ commutes with W. Applying t_ξ to both sides of (4.12) we obtain (assuming $|q| < 1$):

$$\begin{aligned} e^{\hat{\rho}+\beta}\hat{\mathrm{R}}_T = {} & \frac{1}{(n+1)!}\left(\prod_{j=1}^{n+1}\left(\sum_{k_j,r_j=0}^{\infty} - \sum_{k_j,r_j=-1}^{-\infty}\right)\right)(-1)^{\sum_{j=1}^{n+1} r_j} \\ & \times \sum_{k_{n+2},\dots,k_m\in\mathbb{Z}}\sum_{w\in W}\epsilon(w)we^{\hat{\rho}-h^\vee\alpha-\sum_{j=1}^{n+1} r_j\alpha_{2j-1}} \\ & \times q^{\frac{1}{2}h^\vee(\alpha|\alpha)+\sum_{j=1}^{n+1} r_j\left(k_j+\frac{1}{2}\right)-\sum_{j=n+2}^{m} k_j-\frac{1}{4}(n+1)(2m-n)}. \end{aligned} \tag{4.13}$$

Here

$$\hat{\mathrm{R}}_T = \mathrm{R}_0\prod_{n=1}^{\infty}\frac{\prod_{\alpha\in\Delta_0}(1-q^ne^\alpha)}{\prod_{\alpha\in\Delta_1}\left(1+q^{n-\frac{1}{2}}e^\alpha\right)},$$

β is the sum of the $\gamma\in\Delta_{1+}$ such that $(\gamma|\xi)=\frac{1}{2}$,

$$\alpha = \sum_{j=1}^{n+1} k_j\left(\alpha_{2j-1}+\cdots+\alpha_{2n+2}\right) + \sum_{j=n+2}^{m} k_j\alpha_{n+1+j}$$

(so that $(\hat{\rho}|\alpha) = \sum_{j=n+2}^m k_j$). Dividing both sides of (4.13) by R_0 and letting $e^{\alpha_i}\mapsto 1$ ($i = 1,\dots,m+n+1$), we obtain (using the Weyl dimension formula) a power series expansion for the product $\prod_{j=1}^\infty\left(1-q^j\right)^{d_0}/\left(1-q^{j-\frac{1}{2}}\right)^{d_1}$. Furthermore, applying $t_{-\xi}$ to both sides of (4.13), we obtain a better expression for the affine denominator:

$$\begin{aligned} e^{\hat{\rho}}\hat{\mathrm{R}} = {} & \frac{1}{(n+1)!}\sum_{J\subset\{1,\dots,n+1\}}\sum_{\left(\vec{k},\vec{r}\right)\in S_J}\sum_{k_{n+2},\dots,k_m\in\mathbb{Z}}\epsilon_J\left(\vec{k},\vec{r}\right)(-1)^{\sum_{j\in J} r_j} \\ & \times\sum_{w\in W}\epsilon(w)w\frac{e^{\hat{\rho}-h^\vee\alpha-\sum_{j\in J} r_j\alpha_{2j-1}}}{\prod_{j\in J^c}\left(1+e^{-\alpha_{2j-1}}\right)}q^{\frac{1}{2}h^\vee(\alpha|\alpha)+\sum_{j\in J} r_jk_j-\sum_{j=n+2}^m k_j}. \end{aligned} \tag{4.14}$$

Here

$$S_j = \left\{\left(\vec{k},\vec{r}\right)_{j\in J}\in\mathbb{Z}^{2|J|}\,\middle|\, k_j>0 \text{ and } r_j\geq 0,\ \text{or } k_j<0,\ r_j<0 \text{ for each } j\in J\right\},$$

$$\epsilon_J\left(\vec{k},\vec{r}\right) = (-1)^{\sharp\{j\in J|k_j<0\}} \text{ and } J^c = \{1,2,\dots,n+1\}\setminus J.$$

Note that $\left(\rho-h^\vee\alpha-\sum_{j\in J} r_j\alpha_{2j-1}|\alpha_{2i-1}\right)=0$ if $i\in J^c$. Hence, by dividing both sides of (4.14) by R_0 and letting $e^{\alpha_i}\mapsto 1$ ($i=1,\dots,m+n+1$), we may use Theorem 3.2 to obtain a power series expansion for $\prod_{j=1}^\infty\left(1-q^j\right)^{d_0}/\left(1+q^j\right)^{d_1}$.

Example 4.5. Let $\mathfrak{g} = B(m,m)$. We choose a set of simple roots $\alpha_1, \dots, \alpha_{2m} \in \Delta_1$ with the following scalar products: $(\alpha_i|\alpha_i) = 0$ for $i = 1, \dots, 2m-1$, $(\alpha_{2m}|\alpha_{2m}) = -1/2$, $(\alpha_i|\alpha_{i+1}) = (-1)^{i+1}/2$ for $i = 1, \dots, 2m-1$. We have: $h^\vee = 1/2$ and $M^\sharp$ is spanned over $\mathbb{Z}$ by vectors $2\,(\alpha_{2j-1} + \alpha_{2j})$, $j = 1, \dots, m$. Then formula (4.2) can be written using (2.4) and (3.4), as follows:

$$e^{\hat\rho}\hat{\mathrm{R}} = \frac{1}{2^m m!} \sum_{\alpha \in M^\sharp} \sum_{w \in W} \epsilon(w) w t_{-\alpha} \left(e^{\hat\rho} \prod_{j=1}^{m} \left(1 + e^{-\alpha_{2j-1}}\right)^{-1} \right). \tag{4.15}$$

Introduce the element $\xi \in \mathfrak{h}$ by relations $(\xi|\alpha_j) = (-1)^j/2$ for $j = 1, \dots, m-1$, $(\xi|\alpha_{2m}) = 0$. Then $w(\xi) - \xi \in M^\sharp$ for all $w \in W$. Applying t_ξ to both sides of (4.15) we obtain a more convenient for specializations formula:

$$\begin{aligned} e^{\hat\rho+\beta}\hat{\mathrm{R}}_T = {} & \frac{1}{2^m m!} \left(\prod_{j=1}^{m} \left(\sum_{k_j, r_j = 0}^{\infty} - \sum_{k_j, r_j = -1}^{-\infty} \right) \right) (-1)^{\frac{1}{2}m(m+1) + \sum_{j=1}^{m} r_j} \\ & \times \sum_{w \in W} \epsilon(w) w e^{\hat\rho + \frac{1}{2}(\xi - \alpha) - \sum_{j=1}^{m} r_j \alpha_{2j-1}} \\ & \times q^{\frac{1}{4}(\alpha|\alpha) + \sum_{j=1}^{m} \left(k_j(r_j+1) + \frac{1}{2} r_j\right) - \frac{1}{4}m(m-1)}, \end{aligned} \tag{4.16}$$

where $\beta = \frac{1}{2}\xi + \sum_{\gamma \in \Delta_{1+} : (\xi|\gamma) = \frac{1}{2}} \gamma - \sum_{\gamma \in \Delta_{0+} : (\xi|\gamma) = -1} \gamma$ and $\alpha = 2 \sum_{j=1}^{m} k_j \sum_{i=2j-1}^{2m} \alpha_i$.

§5. Applications to sums of squares and sums of triangular numbers.

Consider the following power series in q (that converge for $|q| < 1$):

$$\square(q) = \sum_{n \in \mathbb{Z}} q^{n^2}, \quad \Delta(q) = \sum_{n \in \mathbb{Z}_+} q^{n(n+1)/2}.$$

One has the following product expansions due to Gauss (which follow immediately from the Jacobi triple product identity):

$$\square(-q) = \prod_{k \geq 1} \frac{1 - q^k}{1 + q^k}, \tag{5.1}$$

$$\Delta(q) = \prod_{k \geq 1} \frac{1 - q^{2k}}{1 - q^{2k-1}}. \tag{5.2}$$

Let $\Delta_j = j(j+1)/2$ and let $\Delta = \{\Delta_j | j \in \mathbb{Z}_+\}$ be the set of triangular numbers. Let $\square_{N,k} = |\{\vec{n} \in \mathbb{Z}^N | \sum_{j=1}^{N} n_j^2 = k\}|$ (resp. $\Delta_{N,k} = |\{\vec{n} \in \Delta^N | \sum_{j=1}^{N} n_j = k\}|$) be the number of representations of k as a sum of N squares (resp. triangular numbers). Let $\square_N(q) = \sum_{k\in\mathbb{Z}_+} \square_{N,k}q^k$ and $\Delta_N(q) = \sum_{k\in\mathbb{Z}_+} \Delta_{N,k}q^k$ be the generating series for these numbers. Then, clearly

$$\square_N(q) = \square(q)^N, \quad \Delta_N(q) = \Delta(q)^N.$$

Thus, in view of (5.1) and (5.2) we may use the results of Section 4 to obtain expansion for $\square_N(q)$ and $\Delta_N(q)$. In fact, formulas (4.3) and (4.7) give such expansions provided that $d_0 = d_1$, which happens for $\mathfrak{g}$ of type $A(n, n-1)$, $B(n, n)$ or $D(n+1, n+1)$. These gives us formulas for $\square_N(q)$ and $\Delta_N(q)$, where respectively

$$N = 2n(n+1),\ 2n(2n+1) \text{ or } 4(n+1)^2 \qquad (n \in \mathbb{N}).$$

Example 5.1. $\mathfrak{g} = A(1, 0)$. Letting $z \to 1$ in (4.10) we get:

$$\begin{aligned}\square(-q)^4 &= 1 - 4\sum_{m,n=1}^{\infty} (-1)^{m+n}(m+n)q^{mn} \\ &= 1 + 8\sum_{m,n=1}^{\infty} (-1)^{m+n-1}mq^{mn} \\ &= 1 + 8\sum_{m=1}^{\infty} \frac{m(-q)^m}{1+q^m}.\end{aligned}$$

Since $\frac{2mq^{2m}}{1+q^{2m}} = \frac{2mq^{2m}}{1-q^{2m}} - \frac{4mq^{4m}}{1-q^{4m}}$, we obtain:

$$\square(q)^4 = 1 + 8\sum_{\substack{k\geq 1 \\ 4\nmid k}} \frac{kq^k}{1-q^k},$$

which is Jacobi's formula (0.2). In other words, for $n \geq 1$:

$$\square_{4,n} = 8\sum_{\substack{k\,|\,n \\ 4\nmid k}} k.$$

Letting $z = i$ in (4.9), we get Jacobi's formula (0.1), *i.e.,* that $\square_{2,n}$ equals 4 times the difference between the number of divisors of n congruent to 1 mod 4 and the number of divisors of n congruent to -1 mod 4. This result goes back to Gauss.

Letting in (4.9) $x = -y = q^{1/2}$, we obtain:

$$\Delta(q)^2 = \sum_{j,k=0}^{\infty} (-1)^k q^{((2j+1)(2k+1)-1)/4}.$$

This means that $\Delta_{2,n}$ equals the difference between the number of divisors of $4n+1$ congruent to 1 mod 4 and the number of divisors of $4n+1$ congudent to -1 mod 4. Replacing in (4.9) q by q^2, letting $x = -qz$, $y = -q$ and tending z to 0, we get

$$\Delta(q)^4 = \sum_{j,k=0}^{\infty} (2k+1) q^{((2j+1)(2k+1)-1)/2}.$$

This means that $\Delta_{4,n}$ equals the sum of all divisors of $2n+1$. This result is due to Legendre (1828).

Example 5.2. $\mathfrak{g} = B(1,1)$. Dividing both sides of (4.11) by $(1-xy)(1-y)$ and letting $x = y = z \to 1$, we obtain Jacobi's formula (0.3). In other words,

$$\Box_{6,n} = 4 \sum_{\substack{j,k\geq 1 \\ j\,\mathrm{odd}, jk=n}} (-1)^{\frac{1}{2}(j+1)} (j^2 - 4k^2).$$

Furthermore, replacing in (4.11) q by q^2 and letting $x = y = -qz$, we obtain after dividing both sides by $(z-1)^2$ and letting $z \to 1$:

$$\Delta(q)^6 = -\frac{1}{8} \sum_{\substack{j,k \text{ odd} \\ j>k\geq 1 \\ \frac{1}{2}(j-k)\text{odd}}} (-1)^{\frac{1}{4}(j-1)(k+1)} (j^2 - k^2) q^{\frac{1}{4}(jk-1)}.$$

In other words, as can be easily seen:

$$\Delta_{6,n} = \frac{1}{8} \sum_{\substack{j>k\geq 1 \\ jk=4n+3}} (-1)^{\frac{1}{2}(j+1)} \left(j^2 - k^2\right).$$

Example 5.3. $\mathfrak{g} = A(m, m-1)$. Dividing both sides of (4.13) by R_0, letting $e^{\alpha_i} \to 1$ $(i = 1, \dots, 2m)$ and replacing $q^{\frac{1}{2}}$ by $-q$, we obtain

$$\Delta(q)^{2m(m+1)} = \left(\sum_{\vec{k},\vec{r}\in\mathbb{Z}_+^m} - \sum_{\vec{k},\vec{r}\in(-\mathbb{N})^m} \right) \prod_{j=1}^{m} \frac{r_j' + |\vec{k}|}{(j!)^2} \left(\prod_{1\leq i<j\leq m} (r_i - r_j)(r_i' - r_j') \right)$$
$$\times\, q^{2\sum_{1\leq i\leq j\leq m} k_i k_j + \sum_{j=1}^{m} r_j(2k_j+1) - m(m+1)/2},$$

where

$$r'_j = r_j + k_j. \tag{5.3}$$

Similarly, we obtain from (4.14) using the remarks and notation of Example 4.4:

$$\begin{aligned}\square(-q)^{2m(m+1)} = \frac{(-1)^{\frac{1}{2}m(m+1)}2^m}{\prod_{j=1}^m (j!)^2} & \sum_{J\subset\{1,\dots,m\}} \sum_{(r_j)_{j\in J^c}} \sum_{\left(\vec{k},\vec{r}\right)\in S_J} \epsilon_J\left(\vec{k},\vec{r}\right)(-1)^{\sum_{j=1}^m r_j} \\ & \times 2^{-|J^c|} C\left(\sum_{i\in J^c} r_i + |J^c| - 1,\, m^2 + |J^c| - 1\right) \prod_{j=1}^m \left(r'_j + k_1 + \cdots + k_m\right) \\ & \times \prod_{1\le i<j\le m} \left(r_i - r_j\right)\left(r'_i - r'_j\right) q^{\sum_{1\le i\le j\le m} k_i k_j + \sum_{j\in J} k_j r_j}\end{aligned}$$

Example 5.4. $\mathfrak{g} = B(m,m)$. Dividing both sides of (4.16) by R_0, letting $e^{\alpha_i} \to 1$ $(i = 1, \dots, 2m)$ and replacing $q^{\frac{1}{2}}$ by $-q$ we obtain (using the Weyl dimension formula):

$$\begin{aligned}\Delta(q)^{2m(2m+1)} = \frac{1}{2^m m!} & \left(\sum_{\vec{k},\vec{r}\in\mathbb{Z}_+^m} - \sum_{\vec{k},\vec{r}\in(-\mathbb{N})^m}\right) \prod_{j=1}^m \frac{(-1)^{k_j}}{\left((2j-1)!\right)^2}(1+2r_j)(1+r'_j) \\ & \times \prod_{1\le i<j\le m} \left(r_i - r_j\right)\left(r'_i - r'_j\right)\left(1 + r_i + r_j\right)\left(2 + r'_i + r'_j\right) \\ & \times q^{\sum_{j=1}^m \left(k_j^2 + 2r_jk_j + 2k_j + r_j\right) - \frac{1}{2}m(m-1)}.\end{aligned}$$

Conjecture 5.1.

$$\begin{aligned}\Delta(q)^{2m(2m+1)} = & \frac{1}{m!\left(\prod_{j=1}^m (2j-1)!\right)^2} \sum_{\vec{k},\vec{r}\in\mathbb{Z}_+^m} (-1)^{|\vec{k}|} \\ & \times \prod_{1\le i<j\le m} \left(r_i - r_j\right)\left(r'_i - r'_j\right)\left(2 + r'_i + r'_j\right)\left(1 + r_i + r_j\right) \\ & \times \prod_{j=1}^m \left(1 + 2r_j\right)\left(1 + r'_j\right) q^{\sum_{j=1}^m \left(\left(k_j^2 + 2k_j\right) + \left(r_j(2k_j+1)\right)\right) - \frac{1}{2}m(m-1)}.\end{aligned}$$

§6. Some "degenerate" theta functions and modular forms.

In this section we use the denominator identity (4.8) for $sl(2,1)^\wedge$ to derive transformation properties of certain "degenerate" theta functions and modular forms, and related to them characters of $N = 2$ superconformal algebras.

Let $q = e^{2\pi i\tau}$ where $Im\tau > 0$, and let $\Theta(\tau, z)$ be the standard Jacobi theta function:

$$\begin{aligned}\Theta(\tau, z) &= e^{\pi i\tau/4} e^{-\pi iz} \prod_{n\geq 1} (1-q^n)\left(1-q^n e^{-2\pi iz}\right)\left(1-q^{n-1}e^{2\pi iz}\right)\\ &= \sum_{j\in\mathbb{Z}} (-1)^j q^{\frac{1}{2}(j-\frac{1}{2})^2} e^{2\pi i\left(j-\frac{1}{2}\right)z}.\end{aligned}$$

Recall that this theta function has the following transformation property (see, *e.g.* [K9]):

$$\Theta\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) = i(-i\tau)^{1/2} e^{\pi iz^2/\tau}\Theta(\tau, z).$$

Introduce the following function in τ, z_1, and z_2:

$$F(\tau, z_1, z_2) = \frac{\eta(\tau)^3\Theta(\tau, z_1+z_2)}{\Theta(\tau, z_1)\Theta(\tau, z_2)},$$

where $\eta(\tau)$ is the Dedekind η-function. Note that $F(\tau, z_1, z_2)$ is the left-hand side of identity (4.8) where $x = -e^{2\pi iz_1}$, $y = -e^{2\pi iz_2}$.

Fix a positive integer m. Using the transformation formula for Θ and identity (4.8), it is not hard to derive the following transformation formula:

$$F\left(-\frac{m}{\tau}, \frac{z_1}{\tau}, \frac{z_2}{\tau}\right) = \frac{\tau}{m} e^{\frac{2\pi iz_1z_2}{m\tau}} \sum_{a,b\in\mathbb{Z}/m\mathbb{Z}} e^{\frac{2\pi i(az_1+bz_2)}{m}} q^{\frac{ab}{m}} F(m\tau, z_1+a\tau, z_2+b\tau). \tag{6.1}$$

Given $j, k \in \frac{1}{2}\mathbb{Z}$, introduce the following functions:

$$F^{(m)\pm}_{j,k}(\tau, z) = q^{\frac{jk}{m}} e^{\frac{2\pi i(j-k)z}{m}} F\left(m\tau, -z+j\tau-\tfrac{1}{4}\pm\tfrac{1}{4}, z+k\tau+\tfrac{1}{4}\mp\tfrac{1}{4}\right).$$

We call them *degenerate* theta functions since, due to (4.8), they have the following power series expansion associated to a bilinear form representing zero:

$$F^{(m)\pm}_{j,k}(\tau, z) = \left(\sum_{a,b=0}^{\infty} - \sum_{a,b=-1}^{-\infty}\right) (\pm 1)^{b-a} q^{m\left(a+\frac{k}{m}\right)\left(b+\frac{j}{m}\right)} e^{2\pi i\left(\left(b+\frac{j}{m}\right)-\left(a+\frac{k}{m}\right)\right)z}. \tag{6.2}$$

Transformation formula (6.1) implies the transformation formulas for indefinite theta functions. In order to state them introduce the following sets:

$$A_m^+ = \{(j,k) \in \mathbb{Z}^2 | 0 < j, j+k < m, \quad 0 \le k < m\},$$
$$A_m^- = \left\{(j,k) \in \left(\tfrac{1}{2} + \mathbb{Z}\right)^2 | 0 < j, k, j+k < m\right\}.$$

Introduce the following numbers:

$$S^{(m)}_{(j,k)(a,b)} = \frac{2}{m} e^{\frac{\pi i (j-k)(a-b)}{m}} \sin \frac{\pi (j+k)(a+b)}{m}.$$

Provided that $j, k \in \mathbb{Z}$, one has:

$$F^{(m)\pm}_{j,k}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) = -i\tau e^{-\frac{2\pi i z^2}{m\tau}} \sum_{(a,b)\in A_m^\pm} S^{(m)}_{(j,k)(a,b)} F^{(m)+}_{a,b}(\tau, z). \tag{6.3}$$

Provided that $j, k \in \frac{1}{2} + \mathbb{Z}$, one has:

$$F^{(m)\pm}_{j,k}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) = -i\tau e^{-\frac{2\pi i z^2}{m\tau}} \sum_{(a,b)\in A_m^\pm} S^{(m)}_{(j,k)(a,b)} F^{(m)-}_{a,b}(\tau, z). \tag{6.4}$$

It follows that the linear span of functions $\{F^{(m)+}_{j,k}$ and $F^{(m)-}_{j,k} | (j,k) \in A_m^+ \cup A_m^-\}$ is invariant with respect to the action of $SL_2(\mathbb{Z})$ given by

$$F(\tau, z)|_{\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)} = (c\tau + d)^{-1} e^{\frac{2\pi i c z^2}{m(c\tau+d)}} F\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right).$$

We want now to specialize to $z = 0$. Let $\varphi^{(m)+}_{j,k}(\tau) = F^{(m)+}_{j,k}(\tau, 0)$ for all $(j,k) \in A_m^-$ and $\varphi^{(m)-}_{j,k}(\tau) = F^{(m)-}_{j,k}(\tau, 0)$ for all $(j,k) \in A_m^- \cup A_m^+$ except for $(j, 0) \in A_m^+$ (when the series diverges). These functions have the following power series expansions:

$$\varphi^{(m)\pm}_{j,k}(\tau) = \left(\sum_{a,b=0}^{\infty} - \sum_{a,b=-1}^{-\infty}\right) (\pm 1)^{a+b} q^{m\left(a+\frac{j}{m}\right)\left(b+\frac{k}{m}\right)}.$$

In order to get $SL_2(\mathbb{Z})$-invariance, we must add functions $\varphi^{(m)-}_{j,0}(\tau)$ ($j \in \mathbb{Z}$, $0 < j < m$) given by

$$\varphi^{(m)-}_{j,0} = \tfrac{1}{2} + \left(\sum_{\substack{a,b=0 \\ b>0}}^{\infty} - \sum_{a,b=-1}^{-\infty}\right) (-1)^{a+b} q^{m\left(a+\frac{j}{m}\right)b}.$$

Then we have:

$$\varphi_{j,k}^{(m)+}\left(-\frac{1}{\tau}\right) = -i\tau \sum_{(a,b)\in A_m^+} S_{(j,k)(a,b)}\varphi_{a,b}^{(m)-}(\tau) \text{ if } (j,k)\in A_m^-,$$
$$\varphi_{j,k}^{(m)-}\left(-\frac{1}{\tau}\right) = -i\tau \sum_{(a,b)\in A_m^-} S_{(j,k)(a,b)}\varphi_{a,b}^{(m)-(\text{resp}+)}(\tau) \text{ if } (j,k)\in A_m^{-(\text{resp}+)}.$$

Note also that

$$\varphi_{j,k}^{(m)+}(\tau+1) = e^{\frac{2\pi ijk}{m}}\varphi_{j,k}^{(m)-}(\tau) \text{ if } (j,k)\in A_m^-,$$
$$\varphi_{j,k}^{(m)-}(\tau+1) = e^{\frac{2\pi ijk}{m}}\varphi_{j,k}^{(m)+(\text{resp}-)}(\tau) \text{ if } (j,k)\in A_m^{-(\text{resp}+)}.$$

Thus we have constructed a finite set of modular forms of weight 1 whose linear span is $SL_2(\mathbb{Z})$-invariant.

We derive now, using (6.3) and (6.4) the transformation properties of characters and supercharacters of the $N=2$ superalgebras (*cf.* [RY]). Introduce the following normalized $N=2$ demoninators:

$$\Psi^{(0)\pm}(\tau,z) = e^{\pi iz}\prod_{n=1}^{\infty}\frac{(1-q^n)^2}{(1\pm q^ne^{-2\pi iz})(1\pm q^{n-1}e^{2\pi iz})},$$
$$\Psi^{(1/2)\pm}(\tau,z) = q^{1/8}\prod_{n=1}^{\infty}\frac{(1-q^n)^2}{(1\pm q^{n-1/2}e^{-2\pi iz})(1\pm q^{n-1/2}e^{2\pi iz})}.$$

From the transformation formula for Θ and η one gets:

$$\Psi^{(0)-}\left(-\frac{1}{\tau},\frac{z}{\tau}\right) = \tau e^{-\frac{\pi iz^2}{\tau}}\Psi^{(0)-}(\tau,z),$$
$$\Psi^{(0)+}\left(-\frac{1}{\tau},\frac{z}{\tau}\right) = -i\tau e^{-\frac{\pi iz^2}{\tau}}\Psi^{(1/2)-}(\tau,z),$$
$$\Psi^{(1/2)-}\left(-\frac{1}{\tau},\frac{z}{\tau}\right) = -i\tau e^{-\frac{\pi iz^2}{\tau}}\Psi^{(0)+}(\tau,z),$$
$$\Psi^{(1/2)+}\left(-\frac{1}{\tau},\frac{z}{\tau}\right) = -i\tau e^{-\frac{\pi iz^2}{\tau}}\Psi^{(1/2)+}(\tau,z),$$

One knows that the normalized characters and supercharacters of the unitary discrete series of the two $N=2$ superalgebras corresponding to $\epsilon=0$ or $\frac{1}{2}$ of central charge $3c_m$, where $c_m = 1-\frac{2}{m}$, $m=2,3,\ldots$, are given by the following formulas ([D],[M],[RY]):

$$\mathrm{ch}_{j,k}^{(m,\epsilon)}(\tau,z) = \frac{F_{j,k}^{(m)-}(\tau,z)}{\Psi^{(\epsilon)+}(\tau,z)},$$
$$\mathrm{sch}_{j,k}^{(m,\epsilon)}(\tau,z) = \frac{F_{j,k}^{(m)+}(\tau,z)}{\Psi^{(\epsilon)-}(\tau,z)},$$

if $j, k \in \epsilon + \mathbb{Z}$; the range of (j,k) is A_m^+ or A_m^- depending on whether $\epsilon = 0$ or $\frac{1}{2}$. From the above transformation formulas we obtain the following theorem.

Theorem 6.1. *(cf. [RY]) (a) If $(j,k) \in A_m^-$, then*

$$\begin{aligned}
\mathrm{ch}_{j,k}^{(m,1/2)}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) &= e^{\frac{\pi i z^2}{\tau} c_m} \sum_{a,b \in A_m^-} S_{(j,k)(a,b)}^{(m)} \, \mathrm{ch}_{a,b}^{(m,1/2)}(\tau, z), \\
\mathrm{sch}_{j,k}^{(m,1/2)}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) &= e^{\frac{\pi i z^2}{\tau} c_m} \sum_{a,b \in A_m^+} S_{(j,k)(a,b)}^{(m)} \, \mathrm{ch}_{a,b}^{(m,0)}(\tau, z), \\
\mathrm{ch}_{j,k}^{(m,1/2)}(\tau+1, z) &= e^{\frac{2\pi i jk}{m} - \frac{\pi i}{4}} \, \mathrm{sch}_{j,k}^{(m,1/2)}(\tau, z), \\
\mathrm{sch}_{j,k}^{(m,1/2)}(\tau+1, z) &= e^{\frac{2\pi i jk}{m} - \frac{\pi i}{4}} \, \mathrm{ch}_{j,k}^{(m,1/2)}(\tau, z).
\end{aligned}$$

(b) If $(j,k) \in A_m^+$, then

$$\begin{aligned}
\mathrm{ch}_{j,k}^{(m,0)}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) &= e^{\frac{\pi i z^2}{\tau} c_m} \sum_{a,b \in A_m^-} S_{(j,k)(a,b)}^{(m)} \, \mathrm{sch}_{a,b}^{(m,1/2)}(\tau, z), \\
\mathrm{sch}_{j,k}^{(m,0)}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) &= -i e^{\frac{\pi i z^2}{\tau} c_m} \sum_{a,b \in A_m^+} S_{(j,k)(a,b)}^{(m)} \, \mathrm{sch}_{a,b}^{(m,0)}(\tau, z), \\
\mathrm{ch}_{j,k}^{(m,0)}(\tau+1, z) &= e^{\frac{2\pi i jk}{m}} \, \mathrm{ch}_{j,k}^{(m,0)}(\tau, z), \\
\mathrm{sch}_{j,k}^{(m,0)}(\tau+1, z) &= e^{\frac{2\pi i jk}{m}} \, \mathrm{sch}_{j,k}^{(m,0)}(\tau, z).
\end{aligned}$$

Corollary 6.1. *The linear span of all normalized characters and supercharacters of the minimal unitary discrete series with central charge $3c_m$ is invariant under the action of $SL_2(\mathbb{Z})$ defined by*

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} f(\tau, z) = e^{-\frac{\pi i c z^2}{c\tau+d} c_m} f\left(\frac{a\tau+b}{c\tau+d}, \frac{z}{c\tau+d}\right).$$

Example 6.1. $m = 2$ (minimal possible). Then $A_2^+ = \{(1,0)\}$, $A_2^- = \left\{\left(\frac{1}{2}, \frac{1}{2}\right)\right\}$, and $\mathrm{ch}_{j,k}^{(2,\epsilon)} = \mathrm{sch}_{j,k}^{(2,\epsilon)} = 1$. Furthermore,

$$\begin{aligned}
\varphi_{1/2,1/2}^{+}(\tau) &= \left(F_{1/2,1/2}^{(2)+}(\tau, z) = \Psi^{(1/2)-}(\tau, z)\right)_{z=0} = q^{\frac{1}{8}} \Delta\left(q^{\frac{1}{2}}\right)^2, \\
\varphi_{1/2,1/2}^{-}(\tau) &= \left(F_{1/2,1/2}^{(2)-}(\tau, z) = \Psi^{(1/2)+}(\tau, z)\right)_{z=0} = q^{\frac{1}{8}} \Delta\left(-q^{\frac{1}{2}}\right)^2, \\
\varphi_{1,0}^{-}(\tau) &= \left(F_{1,0}^{(2)-}(\tau, z) = \Psi^{(0)+}(\tau, z)\right)_{z=0} = \tfrac{1}{2}\Box(-q)^2.
\end{aligned}$$

These three functions span a 3-dimensional $SL_2(\mathbb{Z})$-invariant space of modular forms of weight 1. (Note that $F_{1,0}^{(2)+}(\tau, z)$ has a pole at $z = 0$.)

Remark 6.1. Similar results may be obtained for the affine denominator in the general case. As in [K5], this may be used in order to prove the "strange formula" (1.2).

§7. The case $h^\vee = 0$.

The simplest non-trivial case when $h^\vee = 0$ is $\mathfrak{g} = A(1,1)$. It is more convenient to consider $\mathfrak{g} = gl(2,2)$ instead, with simple roots $\alpha_1, \alpha_2, \alpha_3$, such that $(\alpha_1|\alpha_1) = (\alpha_3|\alpha_3) = 0$, $(\alpha_3|\alpha_3) = 2$, $-(\alpha_1|\alpha_2) = (\alpha_3|\alpha_2) = 1$. We conjecture that the denominator identity still holds with an extra factor. Namely, letting $x = e^{-\alpha_1}$, $y = e^{-\alpha_2}$, $z = e^{-\alpha_3}$, introduce the following functions (fixed under $W \ltimes t_{\mathbb{Z}\alpha_2}$):

$$f_n = x^{-n}z^{-n}(x^{2n-1} - z^{2n-1})(x-z) \sum_{j\in\mathbb{Z}_+} (-1)^j q^{(j+1)(j+2n)/2}.$$

Conjecture 7.1. $e^\rho \widehat{\mathrm{R}} = \left(1 + \sum_{n=1}^{\infty} f_n\right) \sum_{n\in\mathbb{Z}} t_{n\alpha_2}(e^\rho \mathrm{R}).$

If we divide both sides by R and let $x, y, z \to 1$ we obtain Jacobi's formula (0.4), *i.e.*,

$$\square_{8,n} = 16 \sum_{j|n} (-1)^{n+j} j^3.$$

So far we have entirely left out the series of simple Lie superalgebras $\mathfrak{g}$ of type $Q(m)$, which do not admit a non-degenerate even invariant bilinear form (but admit an odd one), hence it is natural to let $h^\vee = 0$. Its even part $\mathfrak{g}_0$ is a simple Lie algebra of type A_m and the $\mathfrak{g}_0$-module $\mathfrak{g}_1$ is the adjoint A_m-module. Let Δ (resp Δ_+) denote the set of roots (resp positive roots) of A_m, let $\Pi = \{\alpha_1, \ldots, \alpha_m\}$ be the set of simple roots, let ρ be the half of the sum of roots from Δ_+, let W be the Weyl group, and let $(.|.)$ be the W-invariant bilinear form normalized by $(\alpha_i|\alpha_i) = 2$. Let $s = \left[\frac{m+1}{2}\right]$ and consider the following (maximal orthogonal) set of s roots: $\gamma_1 = \alpha_1 + \ldots + \alpha_m$, $\gamma_2 = \alpha_2 + \ldots + \alpha_{m-1}, \ldots$. The denominator identity for $Q(m)$ is given by the following formula:

$$\mathrm{R} := \prod_{\alpha\in\Delta_+} \frac{1 - e^{-\alpha}}{1 + e^{-\alpha}} = \sum_{w\in W} \epsilon(w) w \prod_{j=1}^{s} (1 + e^{-\gamma_j})^{-1}. \tag{7.1}$$

The associated to $Q(m)$ affine superalgebra $\hat{\mathfrak{g}}$ is a central extension (by $\mathbb{C}C$) of the Lie superalgebra $\bigoplus_{j\in\mathbb{Z}} t^j \mathfrak{g}_{j \bmod 2}$ (its Cartan matrix is not symmetrizable). We let $q=-e^{-C}$ and define the affine denominator as follows:

$$\widehat{\mathrm{R}} = \mathrm{R}\prod_{n=1}^{\infty}\left(\left(\frac{1-q^{2n}}{1-q^{2n-1}}\right)^{2s}\prod_{\alpha\in\Delta}\frac{1-q^{2n}e^{\alpha}}{1-q^{2n-1}e^{\alpha}}\right).$$

In order to state the affine denominator identity, define t_α for $\alpha\in\mathbb{Z}\Delta$ as follows

$$t_\alpha(e^{\beta}) = q^{(\alpha|\beta)}e^{\beta}, \quad \beta\in\mathbb{Z}\Delta, \quad t_\alpha(q)=q.$$

Conjecture 7.2. $$e^{\rho}\widehat{\mathrm{R}} = \sum_{w\in W}\sum_{\vec{n}\in\mathbb{Z}_+^s}\sum_{\substack{\vec{k}\in\mathbb{Z}_+^s\\ k_1\geq\ldots\geq k_s}} \epsilon(w)q^{|\vec{k}|}wt_{n_1\gamma_1+\cdots+n_s\gamma_s}\left(e^{\rho+\sum_i k_i\gamma_i}\right).$$

In particular, for $m=1$ Conjecture 7.2 reads

$$\begin{aligned}\prod_{n=1}^{\infty}\frac{(1-q^{2n})^2(1-q^{2n-2}x)(1-q^{2n}x^{-1})}{(1-q^{2n-1})^2(1-q^{2n-1}x)(1-q^{2n-1}x^{-1})}\\ =\sum_{j,k\in\mathbb{Z}_+} q^{2jk+j+k}(x^{-j}-x^{j+1})\\ =\sum_{n\in\mathbb{Z}}\frac{q^n}{1-q^{2n+1}x},\end{aligned}$$

where $x=e^{-\alpha_1}$. This identity follows from (4.8) by replacing q by q^2, y by $-qx$ and x by $-q$. As we mentioned in Section 5, dividing both sides by $1-x$ and tending x to 1, we recover Legendre's formula $\Delta(q)^4$.

For $m=2$, Conjecture 7.2 reads (here $x=e^{-\alpha_1}$, $y=e^{-\alpha_2}$):

$$\begin{aligned}\prod_{n=1}^{\infty}\frac{(1-q^{2n})^2(1-q^{2n-2}x)(1-q^{2n-2}y)(1-q^{2n-2}xy)}{(1-q^{2n-1})^2(1-q^{2n-1}x)(1-q^{2n-1}y)(1-q^{2n-1}xy)}\\ \times\frac{(1-q^{2n}x^{-1})(1-q^{2n}y^{-1})(1-q^{2n}x^{-1}y^{-1})}{(1-q^{2n-1}x^{-1})(1-q^{2n-1}y^{-1})(1-q^{2n-1}x^{-1}y^{-1})}\\ =\sum_{j\in\mathbb{N}}\sum_{k\in\mathbb{N}\mathrm{odd}} q^{jk-1}(xy)^{1-j}(1-x^j)(1-y^j)(1-x^jy^j).\end{aligned}$$

Letting in this identity $x=y=z$, we obtain after dividing by $(1-z)^2(1-z^2)$:

$$\begin{aligned}\prod_{n=1}^{\infty}\frac{(1-q^{2n})^2(1-q^{2n}z)^2(1-q^{2n}z^{-1})^2(1-q^{2n}z^2)(1-q^{2n}z^{-2})}{(1-q^{2n-1})^2(1-q^{2n-1}z)^2(1-q^{2n-1}z^{-1})^2(1-q^{2n-1}z^2)(1-q^{2n-1}z^{-2})}\\ =\sum_{j\in\mathbb{N}}\sum_{k\in\mathbb{N}\,\mathrm{odd}} q^{jk-1}z^{2-2j}\left(\frac{z^j-1}{z-1}\right)^2\frac{z^{2j}-1}{z^2-1}.\end{aligned}$$

Letting z tend to 1, we obtain another Legendre's formula:

$$\Delta(q)^8 = \sum_{j\in\mathbb{N}} \sum_{k\in\mathbb{N}\,\mathrm{odd}} j^3 q^{jk-1}, \qquad i.e., \quad \Delta_{8,n} = \sum_{\substack{k|n+1\\ k\,\mathrm{odd}}} \left(\frac{n+1}{k}\right)^3 .$$

Finally, dividing both sides of the identity for general m by R and evaluating at 0 using Weyl's character formula we obtain the following two beautiful results for $m = 2s-1$ (resp. $2s$):

$$\text{(7.2)} \qquad \Delta_{4s^2,n} \left(\mathrm{resp}\, \Delta_{4s(s+1),n}\right) = \sum_{\vec{r}\in\mathbb{Z}_+^s} \sum_{\vec{k}\in F_{\vec{r},n}} \dim L\left(k_1\gamma_1 + \cdots + k_s\gamma_s, A_m\right),$$

where $F_{\vec{r},n} = \left\{\vec{k} \in \mathbb{Z}_+^s | k_1 \geq k_2 \geq \ldots \geq k_s,\ \sum_{i=1}^s (k_i(2r_i+1) + (m+2-2i)r_i) = n\right\}$, and $L(\Lambda, A_m)$ denote the irreducible (finite-dimensional) module over A_m with highest weight Λ.

The simplest new formula corresponds to $m = 3$:

$$\Delta_{16,n} = \frac{1}{12} \sum_{\substack{r_1,r_2\in\mathbb{Z}_+\\ k_1\geq k_2\in\mathbb{Z}_+\\ k_1(2r_1+1)+k_2(2r_1+1)+3r_1+r_2=n}} (2k_1+3)(2k_2+1)(k_1-k_2+1)^2(k_1+k_2+2)^2.$$

Letting $a = 2k_1+3$, $b = 2k_2+1$, $r = 2r_1+1$, $s = 2r_2+1$, we get the following beautiful form of this formula:

$$\Delta_{16,n} = \frac{1}{3\cdot 4^3} \sum_{\substack{a,b,r,s\in\mathbb{N}\mathrm{odd}\\ a>b\\ ar+bs=2n+4}} ab\left(a^2-b^2\right)^2 .$$

Next new formula corresponds to $m = 4$:

$$\Delta_{24,n} = \frac{1}{72} \sum_{\substack{a,b\in\mathbb{N}\\ r,s\in\mathbb{N}\mathrm{odd}\\ a>b\\ ar+bs=n+3}} a^3b^3\left(a^2-b^2\right)^2 .$$

In a similar way the general formula (7.2) can be rewritten as follows:

$$\Delta_{4s^2,n} = \frac{4^{-s(s-1)}}{\prod_{j=1}^{2s-1} j!} \sum_{\substack{a_1,\ldots,a_s\in\mathbb{N}\mathrm{odd}\\ r_1,\ldots,r_s\in\mathbb{N}\mathrm{odd}\\ a_1>\ldots>a_s\\ a_1r_1+\cdots+a_sr_s=2n+s^2}} a_1\ldots a_s \prod_{i<j}\left(a_i^2-a_j^2\right)^2$$

$$\Delta_{4s(s+1),n} = \frac{2^s}{\prod_{j=1}^{2s} j!} \sum_{\substack{a_1,\ldots,a_s\in\mathbb{N}\\ r_1,\ldots,r_s\in\mathbb{N}\mathrm{odd}\\ a_1>\ldots>a_s\\ a_1r_1+\cdots+a_sr_s=n+\frac{1}{2}s(s+1)}} (a_1\ldots a_s)^3 \prod_{i<j}\left(a_i^2-a_j^2\right)^2 .$$

§8. Integrable highest weight modules over $\hat{\mathfrak{g}}$.

Given a choice of $\Delta_+ \subset \Delta$, we associate to each $\Lambda \in \hat{\mathfrak{h}}^*$ an *irreducible highest weight module* $\hat{L}(\Lambda)$ over $\hat{\mathfrak{g}}$ defined by the property that there exists a non-zero vector v_Λ such that:

$$h(v_\Lambda) = \Lambda(h)v_\Lambda \text{ for } h \in \hat{\mathfrak{h}}, \quad \mathfrak{n}_+(v_\Lambda) = 0, \quad (t^n \otimes \mathfrak{g})v_\Lambda = 0 \text{ for } n > 0.$$

The number $c = \Lambda(C)$ is called the *level* of $\hat{L}(\Lambda)$. Note that $L(\Lambda) := U(\mathfrak{g})v_\Lambda$ is an irreducible highest weight module over $\mathfrak{g}$.

Definition 8.1. *A $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is called integrable if the following two properties hold:*

(i) $\dim L(\Lambda) < \infty$,

(ii) $t^n \otimes \mathfrak{g}_\alpha$ are locally nilpotent for all $\alpha \in \Delta_0^\sharp$ and $n \in \mathbb{Z}$.

It is clear that if $\hat{L}(\Lambda)$ is an integrable highest weight $\hat{\mathfrak{g}}$-module for some choice of Δ_+, then it is an integrable highest weight module for any other choice of Δ_+; also, $\hat{L}(\Lambda)$ is $\mathfrak{g}$-locally finite. The $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is integrable if and only if $\mathrm{ch}\, \hat{L}(\Lambda)$ is invariant with respect to $W \ltimes t_{M^\sharp}$.

In order to classify integrable highest weight $\hat{\mathfrak{g}}$-modules $\hat{L}(\Lambda)$, choose Δ_+ such that Π has a unique odd root (*cf.* Example 3.3). Note that α_0 is even if and only if $\mathfrak{g}$ is of the second type; in this case define $\tilde{\theta} \in \Delta_+ \subset \hat{\Delta}_+$ as in Example 3.1c. If $\mathfrak{g}$ is of the first kind, *i.e.,* of type $A(m,n)$ or $C(n)$, we define $\tilde{\theta} = \alpha_0 + \tilde{\theta}'$, where the coefficients of $\tilde{\theta}'$ in terms of Π are the following:

$$\begin{array}{cl} A(m,n),\ m \geq n & \bigcirc - \bigcirc \cdots - \bigcirc - \bigotimes - \bigcirc_1 - \ldots - \bigcirc_1 \\ C(m),\ m > 2 & \bigotimes^1 - \bigcirc - \bigcirc - \cdots - \bigcirc \Leftarrow \bigcirc. \end{array}$$

As before, we let $k = (\Lambda|\tilde{\theta}^\vee)$ and $k_i = (\Lambda|\alpha_i^\vee)$ for all non-isotropic α_i.

Theorem 8.1. *(a) For $\mathfrak{g}$ of the second kind, the $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is integrable if and only if* $\dim L(\Lambda) < \infty$ *(the corresponding conditions on k and the k_i, $i \neq 0$, are given in [K2, p. 84]) and $k_0 \in \mathbb{Z}_+$.*

(b) For $\mathfrak{g}$ of type $A(m,n)$, $m \geq n$, the $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is integrable if and only if

(i) $k \in \mathbb{Z}_+$ and $k_i \in \mathbb{Z}_+$ for $i \neq 0$ and $i \neq m+1$;

(ii) when $k \leq n$, there exist $r, s \in \mathbb{Z}_+$ such that: $r + s = k$, $(\Lambda|\alpha_0) = r + k_{m+n+1} + k_{m+n} + \cdots + k_{m+n-r+2}$, $(\Lambda|\alpha_{m+1}) = s + k_{m+2} + k_{m+3} + \cdots + k_{m+s+1}$.

The $\hat{C}(m)$-module $\hat{L}(\Lambda)$ is integrable if and only if

(i) $k \in \mathbb{Z}_+$ and $k_i \in \mathbb{Z}_+$ for $i \neq 0, 1$,

(ii) if $k = 0$, then $(\Lambda|\alpha_0) = (\Lambda|\alpha_1) = 0$.

Proof. We use two facts: (i) if α is a simple even root, then the root vector attached to $-\alpha$ is locally nilpotent on $\hat{L}(\Lambda)$ if and only if $(\Lambda|\alpha^\vee) \in \mathbb{Z}_+$; (ii) any root from $\widetilde{\Delta}$ can be made simple by applying a sequence of odd simple reflections. ■

One has the following expressions for the affine central charge c:

$A(m,n),\ m \geq n :$ $c = k_1 + \cdots + k_m + k.$
$C(m) :$ $c = k_2 + \cdots + k_m + k.$
$B(m,n),\ D(m,n) :$ $c = k_0 + k_1 + \ldots + k_{n-1} + k.$
$D(2,1;\alpha),\ F(4),\ G(3) :$ $c = k_0 + k.$

It follows that the affine central charge c of an integrable $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is a non-negative integer and that $c = 0$ if and only if the $\hat{\mathfrak{g}}$-module $\hat{L}(\Lambda)$ is 1-dimensional (which happens if and only if $\Lambda|_{\mathfrak{h}+\mathbb{C}C} = 0$).

Example 8.1. $\hat{\mathfrak{g}} = \hat{A}(1,0)$. We choose $\Pi = \{\alpha_1, \alpha_2\}$ where both α_1 and α_2 are odd. Then $(\alpha_1|\alpha_1) = (\alpha_2|\alpha_2) = 0$, $(\alpha_1|\alpha_2) = 1$. Then $\alpha_0 = C - \alpha_1 - \alpha_2$ is an even root. Let $m_i = (\Lambda|\alpha_i)$, $i = 0, 1, 2$. One has $\dim L(\Lambda) < \infty$ if and only if $m_1 + m_2 \in \mathbb{N}$ or $m_1 = m_2 = 0$, and $\hat{L}(\Lambda)$ is integrable if, in addition, $m_0 \in \mathbb{Z}_+$. The central charge $c = m_0 + m_1 + m_2$ $(\in \mathbb{Z}_+)$. Assume that the following "non-degeneracy" condition holds:

$$m_i \neq 0 \ (i = 1 \text{ or } 2) \Rightarrow m_i \text{ is not divisible by } c + 1.$$

Then the same argument as in Section 2 gives the following character formula:

$$e^{\hat{\rho}} \widehat{\mathrm{R}} \operatorname{ch} \hat{L}(\Lambda) = \sum_{\alpha \in M^\sharp} t_\alpha \left(e^{\hat{\rho}} \mathrm{R} \operatorname{ch} L(\Lambda) \right). \tag{8.1}$$

In particular, if $m_1 = m_2 = 0$, we obtain for $c \in \mathbb{Z}_+$:

$$e^{\hat{\rho}} \widehat{\mathrm{R}} \operatorname{ch} \hat{L}(cd) = \sum_{\alpha \in M^\sharp} t_\alpha \left(e^{\hat{\rho} + cd} \mathrm{R} \right). \tag{8.2}$$

Letting $x = e^{-\alpha_1}$, $y = e^{-\alpha_2}$ we may rewrite (8.2) as follows:

$$e^{-cd} \widehat{\mathrm{R}} \operatorname{ch} \hat{L}(cd) = \sum_{\substack{j,k \in \mathbb{Z}_+ \\ \min(j,k) \in (c+1)\mathbb{Z}}} (-1)^{j+k} x^j y^k q^{\frac{jk}{c+1}} - \sum_{\substack{j,k \in \mathbb{N} \\ \min(j,k) \in (c+1)\mathbb{N}}} (-1)^{j+k} x^{-j} y^{-k} q^{\frac{jk}{c+1}}. \tag{8.3}$$

Definition 8.2. *An integrable* $\hat{\mathfrak{g}}$*-module* $\hat{L}(\Lambda)$ *is called* tame *if its character is given by formula (8.1).*

Theorem 4.1 says that the 1-dimensional $\hat{\mathfrak{g}}$-module is tame.

References

[BL] I. N. Bernstein, D.A. Leites, Character formulae for irreducible representations of Lie superalgebras of series *gl* and *sl*, *C.R. Acad. Bulg. Sci* **33** (1980), 1049–51.

[D] V.K. Dobrev, Characters of the unitarizable highest weight modules over the $N = 2$ superconformal algebras, *Phys. Lett.* **186B** (1987), 43–51.

[F] N.J. Fine, *Some basic hypergeometric series and applications*, Math. Surveys **27**, 1988, AMS, Providence.

[H] D. Hickerson, A proof of the mock theta conjectures, *Invent. Math.* **94** (1988), 639–660.

[J] C.G.J. Jacobi, Fundamenta nova theoriae functionum ellipticarum, *Crelle J.* (1829), 55–239.

[JHKT1] J. Van der Jeugt, J.W.B. Hughes, R.C. King and J. Thierry-Mieg, Character formulae for irreducible modules of the Lie superalgebra $sl(m/n)$, *J. Math. Phys.* **31** (1990), 2278–.

[JHKT2] ——, A character formula for singly atypical modules of the Lie superalgebra $sl(m/n)$, *Comm. Alg.* **18** (1990), 3453–3480. .

[K1] V.G. Kac, Infinite-dimensional Lie algebras and Dedekind's η-function, *Funct. Analis i ego Prilozh.* **8** (1974), No. 1, 77–78. English translation: *Funct. Anal. Appl.* **8** (1974), 68–70.

[K2] ——, Lie superalgebras, *Advances in Math.* **26**, No. 1 (1977), 8–96.

[K3] ——, Characters of typical representations of classical Lie superalgebras, *Comm. Alg.* **5** (1977), 889–897.

[K4] ——, Representations of classical Lie superalgebras, *Lect. Notes Math.* **676**, Springer-Verlag (1978), 597–626.

[K5] ——, Infinite-dimensional algebras, Dedekind's η-function, classical Möbious function and the very strange formula, *Advances in Math.* **30** (1978), 85–136.

[K6] ——, Contravariant form for infinite-dimensional Lie algebras and superalgebras, in *Lecture Notes in Physics* **94**, Springer-Verlag (1979) 441–445.

[K7] ——, Laplace operators of infinite-dimensional Lie algebras and theta functions, *Proc. Natl. Acad. Sci. USA* **81** (1984), 645–647.

[K8] ——, Highest weight representations of conformal current algebras, Symposium on Topological and Geometrical methods in Field theory, Espoo, Finland, 1986. World. Sci., 1986, 3–16.

[K9] ——, Infinite-dimensional Lie algrebras, 3rd edition, Cambridge University Press, 1990.

[KL] V.G. Kac, J. van de Leur, Super boson-fermion correspondence, *Ann. d'Institute Fourier* **37** (1987), 99–137.

[KP] V.G. Kac, D.H. Peterson, Infinite-dimensional Lie algebras, theta functions and modular forms, *Advances in Math.* **53** (1984) 125–264.

[Ko] B. Kostant, On Macdonald's η-function formula, the Laplacian and generalized exponents, *Advances in Math.* **20** (1976) 179–212.

[Mac] I.G. Macdonald, Affine root systems and Dedekind's η-function, *Invent. Math.* **15** (1972) 91–143.

[M] Y. Matsuo, Character formula of $c < 1$ unitary representation of $N = 2$ superconformal algebra, *Prog. Theor. Phys.* **77** (1987), 793–797.

[P] I. Penkov, Generic representation of classical Lie superalgebras and their localization, Univ. Cal. Riverside preprint (1992).

[PS1] I. Penkov, V. Serganova, Cohomology of G/P for classical Lie supergroups and characters of some atypical G-modules, *Ann. Inst. Fourier* **39** (1989), 845–873.

[PS2] ——, Representations of classical Lie superalgebras of type I, *Indag. Math* **N.S.3(4)** (1992), 419–466.

[RY] F. Ravanini, S.-K. Yang, Modular invariance in $N = 2$ superconformal field theories, *Phys. Lett.* **195B** (1987), 202–208.

[S] V. Serganova, Kazhdan-Lusztig polynomials for the Lie superalgebras $GL(m/n)$ Yale Univ. preprint (1992).

[Se] A. Sergeev, Tensor algebra of the defining representation as the module over Lie superalgebras $gl(m,n)$ and $Q(n)$, *Math. USSR, Sbornik* **51** (1985), 419–424.

[TM1] J. Thierry-Mieg, Table des representations irreducibles des superalgebras de Lie, unpublished (1983).

[TM2] ——, Irreducible representations of the basic classical Lie superalgebras $SU(m/n)$, $SU(n/n)/\,U(1)$, $OSp(m/2n)$, $D(2/1,\alpha)$, $G(3)$ and $F(4)$, *Lect. Notes in Physics* **201**, eds. G. Denardo, G. Ghirardi and T. Weber, Springer, Berlin, 94–98.

[VJ1] J. Van der Jeugt, Irreducible representations of the exceptional Lie superalgebras $\mathbf{D}(2,1:\alpha)$, *J. Math. Phys.* **26** (1985), 913–924.

[VJ2] ——, Character formulae for the Lie superalgebra $C(n)$, *Comm. Alg.*